

Datasheet: MCA4772F

Description:	RAT ANTI MOUSE TREM-2:FITC
Specificity:	TREM-2
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	78.18
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			
Immunohistology - Frozen			■	
Immunohistology - Paraffin			■	
ELISA			■	
Immunoprecipitation			■	
Western Blotting			■	
Immunofluorescence			■	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Mouse		
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide (NaN ₃)		
Stabilisers	1% Bovine Serum Albumin		
Approx. Protein Concentrations	IgG concentration 0.1mg/ml		
Immunogen	Mouse TREM-2 Fc fusion protein.		

**External Database
Links**

UniProt:

[Q99NH8](#)

[Related reagents](#)

Entrez Gene:

[83433](#)

Trem2

[Related reagents](#)

Synonyms

Trem2a, Trem2b, Trem2c

Fusion Partners

Lymph node cells from immunised Fisher rats were fused with cells of the rat YB-2/0 myeloma cell line.

Specificity

Rat anti Mouse Trem-2 antibody, clone 78.18 recognizes Triggering receptor expressed on myeloid cells 2 (TREM-2), an activating receptor of the Ig superfamily. TREM-2 forms a receptor signalling complex with DAP12 to trigger the activation of the immune response in macrophages and dendritic cells. It may be involved in chronic inflammation and may promote the production of constitutive rather than inflammatory chemokines and cytokines. TREM-2 may also have a functional role in osteoclast formation.

Stimulation of TREM-2 with Rat anti Mouse Trem-2 antibody, clone 78.18 enhances multinucleated osteoclast formation *in vitro*. Rat anti Mouse Trem-2 antibody, clone 78.18 may also be used to block TREM-2 on pre-osteoclasts to inhibit the formation of multinuclear osteoclasts and to inhibit bone resorption by mature osteoclasts.

Flow Cytometry

Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul.

References

1. Humphrey, M.B. *et al.* (2006) TREM2, a DAP12-associated receptor, regulates osteoclast differentiation and function. [J Bone Miner Res. 21 \(2\): 237-45.](#)

Storage

Store at +4°C or at -20°C if preferred.

Storage in frost-free freezers is not recommended.

This product should be stored undiluted. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life

18 months from date of despatch.

Acknowledgements

This product is for *in vitro* research use only. It is not to be used for commercial purposes. Use of this product to produce products for sale or for diagnostic, therapeutic or drug discovery purposes is prohibited. In order to obtain a license to use this product for commercial purposes, contact The Regents of the University of California.

**Health And Safety
Information**

Material Safety Datasheet documentation #10041 available at:
10041: <https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf>

Regulatory

For research purposes only

Related Products

Recommended Negative Controls

[RAT IgG1 NEGATIVE CONTROL:FITC \(MCA1211F\)](#)

Recommended Useful Reagents

[MOUSE SEROBLOCK FcR \(BUF041A\)](#)

[MOUSE SEROBLOCK FcR \(BUF041B\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M302143:170109'

Printed on 05 May 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)