

Datasheet: MCA5776GA

Description:	MOUSE ANTI HUMAN ACTIN GAMMA
Specificity:	ACTIN GAMMA
Other names:	ACTG1
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	2A3
Isotype:	IgG2b
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			■	
Immunohistology - Frozen			■	
Immunohistology - Paraffin	■			
ELISA			■	
Western Blotting	■			1/100 - 1/1000
Immunofluorescence	■			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human
Species Cross Reactivity	Reacts with: Mouse Based on sequence similarity, is expected to react with: Mammals N.B. Antibody reactivity and working conditions may vary between species.
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Carrier Free	Yes
Approx. Protein	IgG concentration 1.0 mg/ml

Concentrations

Immunogen Synthetic peptide corresponding to the N-terminal nonapeptide of gamma-cytoplasmic actin, coupled to KLH.

External Database Links

UniProt:

[P63261](#) [Related reagents](#)

Entrez Gene:

[71](#) ACTG1 [Related reagents](#)

Synonyms ACTB, ACTG

Specificity **Mouse anti Human actin gamma antibody, clone 2A3** recognises cytoplasmic actin gamma (known also as ACTG and Gamma Actin), a 41.8 kDa cytoskeletal protein.

Actins are highly conserved proteins that are involved in cell motility, structure and integrity and are ubiquitously expressed in all eukaryotic cells. Six different isoforms of actin have been identified ([Vandekerckhove and Weber 1978](#)).

Clone 2A3 is highly specific for actin gamma and does not react with other actin isoforms ([Dugina et al. 2009](#)).

References

1. Dugina, V. *et al.* (2009) Beta and gamma-cytoplasmic actins display distinct distribution and functional diversity. [J Cell Sci. 122: 2980-8.](#)
2. Brockmann, C. *et al.* (2011) Beta- and gamma-cytoplasmic actins are required for meiosis in mouse oocytes. [Biol Reprod. 85: 1025-39.](#)
3. Miazza, V. *et al.* (2011) Sendai virus induced cytoplasmic actin remodeling correlates with efficient virus particle production. [Virology. 410: 7-16.](#)
4. Baranwal, S. *et al.* (2012) Nonredundant roles of cytoplasmic β - and γ -actin isoforms in regulation of epithelial apical junctions. [Mol Biol Cell. 23: 3542-53.](#)
5. Dugina, V. *et al.* (2015) Tumor promotion by γ and suppression by β non-muscle actin isoforms. [Oncotarget. 6 \(16\): 14556-71.](#)
6. Simiczyjew A *et al.* (2015) Active invadopodia of mesenchymally migrating cancer cells contain both β and γ cytoplasmic actin isoforms. [Exp Cell Res. pii: S0014-4827\(15\)30146-4.](#)
7. Simiczyjew, A. *et al.* (2017) Involvement of β - and γ -actin isoforms in actin cytoskeleton organization and migration abilities of bleb-forming human colon cancer cells. [PLoS One. 12 \(3\): e0173709.](#)
8. Chaponnier, C. & Gabbiani, G. (2016) Monoclonal antibodies against muscle actin isoforms: epitope identification and analysis of isoform expression by immunoblot and immunostaining in normal and regenerating skeletal muscle. [F1000Res. 5: 416.](#)
9. Chen, A. *et al.* (2017) Cytokinesis requires localized β -actin filament production by an actin isoform specific nucleator. [Nat Commun. 8 \(1\): 1530.](#)

Storage Store at +4°C or at -20°C if preferred.
Storage in frost-free freezers is not recommended.
This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life 18 months from date of despatch.

**Health And Safety
Information**

Material Safety Datasheet documentation #10040 available at:
10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87...)	Alk. Phos. , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (STAR77...)	HRP
Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Rabbit Anti Mouse IgG (STAR8...)	DyLight®800
Goat Anti Mouse IgG (STAR70...)	FITC
Rabbit Anti Mouse IgG (STAR13...)	HRP
Human Anti Mouse IgG2b (HCA038...)	FITC , HRP
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®549 , DyLight®649 , DyLight®680 , DyLight®800 , FITC , HRP

Recommended Negative Controls

[MOUSE IgG2b NEGATIVE CONTROL \(MCA691\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
--------------------------------------	---	------------------	---	---------------	---

'M318628:180718'

Printed on 01 Aug 2018