

Datasheet: MCA5938B

Description:	RAT ANTI MOUSE CD335:Biotin
Specificity:	CD335
Other names:	NKp46
Format:	Biotin
Product Type:	Monoclonal Antibody
Clone:	29A1.4
Isotype:	IgG2a
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			1/5 - 1/10
Immunohistology - Frozen			■	
Immunohistology - Paraffin			■	
ELISA			■	
Western Blotting			■	
Immunofluorescence			■	
Functional Assays (1)			■	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1) **Bio-Rad recommends the use of [MCA5938EL](#) for functional assays.**

Target Species	Mouse
Product Form	Purified IgG conjugated to Biotin - liquid
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Approx. Protein Concentrations	IgG concentration 0.5 mg/ml
Immunogen	NKP46-IgG1 Fc fusion protein.
External Database Links	UniProt:

Entrez Gene:

[17086](#) Ncr1 [Related reagents](#)

Synonyms	Ly94
Specificity	Rat anti Mouse CD335 antibody, clone 29A1.4 recognizes natural killer cell p46-related protein (NKp46), otherwise known as CD335, which is uniquely expressed by resting and activated natural killer (NK) cells, but no expression of CD335 has been detected on B cells, T cells, monocytes or granulocytes. It is a major NK cell lysis receptor for autologous pathogen-infected and tumor target cells during natural cytotoxicity responses. Ligands of CD335 include viral hemagglutinins (HAs) and heparan sulfate proteoglycans (HSPGs) on the surface of tumor cells. Staining with Rat anti mouse CD335 (29A1.4) is not strain specific and the antibody has been used to stain C57BL/6, SJL, CBA/CA and BALB/c strains. Clone 29A1.4 also activates NK cells <i>in vitro</i>. It does not deplete NK cells <i>in vivo</i>.
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul.
References	1. Walzer, T. <i>et al.</i> (2007) Identification, activation, and selective <i>in vivo</i> ablation of mouse NK cells via NKp46. Proc Natl Acad Sci U S A. 104: 3384-9. 2. Walzer, T. <i>et al.</i> (2007) Natural killer cell trafficking <i>in vivo</i> requires a dedicated sphingosine 1-phosphate receptor. Nat Immunol. 8 (12): 1337-44. 3. Luci, C. <i>et al.</i> (2009) Influence of the transcription factor RORgammat on the development of NKp46+ cell populations in gut and skin. Nat Immunol. 10: 75-82. 4. Chiossone, L. <i>et al.</i> (2009) Maturation of mouse NK cells is a 4-stage developmental program. Blood. 113: 5488-96.
Storage	Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Shelf Life	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation available at: Material Safety Datasheet Documentation #10040 available at: https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf
Regulatory	For research purposes only

Related Products

Recommended Useful Reagents

[MOUSE SEROBLOCK FcR \(BUF041A\)](#)

[MOUSE SEROBLOCK FcR \(BUF041B\)](#)

North & South America Tel: +1 800 265 7376
Fax: +1 919 878 3751

Worldwide Tel: +44 (0)1865 852 700
Fax: +44 (0)1865 852 739

Europe Tel: +49 (0) 89 8090 95 21
Fax: +49 (0) 89 8090 95 50

'M264959:150107'

Printed on 04 Oct 2017