

Datasheet: MCA5973PE

Description:	MOUSE ANTI PIG CD27:RPE
Specificity:	CD27
Other names:	SWC2
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	B30C7
Isotype:	IgG1
Quantity:	100 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Pig		
Product Form	Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized		
Reconstitution	Reconstitute with 1ml distilled water		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	RPE 488nm laser	496	578
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)		
	1% Bovine Serum Albumin		
	5% Sucrose		
Immunogen	Porcine peripheral blood monocytes.		
External Database Links	UniProt:		
	F1SL30	Related reagents	

Fusion Partners	Spleen cells from immunized Balb/c mice were fused with cells of the SP2/0-Ag14 myeloma cell line
Specificity	Mouse anti Pig CD27 antibody, clone B30C7 recognizes the porcine homologue of human CD27, previously known as Swine Workshop Cluster 2 (SWC2), a T-cell co-stimulatory molecule belonging to the TNF receptor family. In humans the CD27 antigen is expressed by discrete populations of T- and B-cells where it functions in a co-stimulatory role to induce proliferation of T-cells and B-cells, however studies undertaken using the B30C7 clone have been unable to detect expression of CD27 on porcine B-cells (Reutner et al. 2012). Studies in humans have demonstrated CD27 to be involved in apoptosis, research in swine has shown that porcine CD27 is expressed by all naïve CD8a^{-ve} T-helper cells and a sub-population of CD8a^{+ve} cells (Reutner et al. 2012). Monoclonal antibodies to CD27 have previously been used to differentiate between subsets of NK cells and research utilizing clone B30C7 has shown that this monoclonal antibody may be used to differentiate between subsets of pig NK cells (Mair et al. 2013).
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul
References	1. Reutner, K. <i>et al.</i> (2012) Porcine CD27: identification, expression and functional aspects in lymphocyte subsets in swine. Dev Comp Immunol. 38: 321-31. 2. Reutner, K. <i>et al.</i> (2013) CD27 expression discriminates porcine T helper cells with functionally distinct properties. Vet Res. 44: 18. 3. Franzoni, G. <i>et al.</i> (2013) Assessment of the Phenotype and Functionality of Porcine CD8 T Cell Responses following Vaccination with Live Attenuated Classical Swine Fever Virus (CSFV) and Virulent CSFV Challenge. Clin Vaccine Immunol. 20: 1604-16. 4. Mair, K.H. <i>et al.</i> (2013) Porcine CD8αdim/-NKp46high NK cells are in a highly activated state. Vet Res. 44: 13.
Storage	Prior to reconstitution store at +4°C. Following reconstitution store at +4°C. DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.
Shelf Life	12 months from date of reconstitution
Health And Safety Information	Material Safety Datasheet documentation #10075 available at: 10075: https://www.bio-rad-antibodies.com/uploads/MSDS/10075.pdf
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:RPE \(MCA928PE\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

Printed on 05 May 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)