

Datasheet: MCA6035

Description:	MOUSE ANTI MONO- AND POLYUBIQUITIN
Specificity:	MONO- AND POLYUBIQUITIN
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	FK2
Isotype:	IgG1
Quantity:	50 µg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
ELISA	■			
Immunoprecipitation	■			
Western Blotting	■			1/100 - 1/1000
Immunocytochemistry	■			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Broad
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by ion exchange chromatography from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.1% Sodium Azide (NaN ₃)
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Poly-ubiquitinated lysozyme

Specificity **Mouse anti mono- and polyubiquitin antibody**, clone FK2 recognizes both mono- and polyubiquitin chains. The antibody does not recognize free ubiquitin.

Ubiquitin, as the name implies, is a ubiquitously expressed and highly conserved protein of 8.6 kDa. The protein is covalently linked to selected lysine residues in a post-translational modification

process known as ubiquitylation or ubiquitination. This chemical reaction is mediated by three different protein families; ubiquitin-activating enzymes (also known as E1s), ubiquitin-conjugating enzymes (also known as E2s) and ubiquitin ligases (also known as E3s) ([Hershko and Ciechanover 1998](#)).

The impact of ubiquitination depends on whether a single ubiquitin moiety (monoubiquitination) or an ubiquitin chain (polyubiquitination) has been attached to a protein. Monoubiquitination tends to trigger cellular processes related to endocytosis and membrane trafficking ([Haglund *et al.* 2003](#)) while the impact of polyubiquitination varies depending on how the ubiquitin residues in the chain have been linked. Attachment of Lysine-48 ubiquitin chains results in degradation by the 20S proteasome while addition of Lysine-63 ubiquitin chains mediates DNA damage and NFkappaB signaling ([Chen 2005](#); [Mocciaro and Rape 2012](#)). Lysine-6, Lysine-11, Lysine-27, Lysine-29 and Lysine-33 chains have also been reported ([Komander 2009](#); [Ye and Rape 2011](#)).

When comparing staining of mouse anti-mono and polyubiquitin antibody (clone FK2) against staining with mouse anti polyubiquitin antibody (clone FK1) one can determine if a protein target is mono- or polyubiquitinated.

Western Blotting	Mouse anti mono- and poly-ubiquitin recognizes mono- and poly-ubiquitinated protein conjugates but does not recognize free ubiquitin by Western Blot. Use of milk based blocking reagents is not recommended. 1% BSA in PBS or TBS Tween should be used instead.
Storage	Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Shelf Life	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10303 available at: 10303: https://www.bio-rad-antibodies.com/uploads/MSDS/10303.pdf
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87...)	Alk. Phos. , HRP
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®549 , DyLight®649 , DyLight®680 , DyLight®800 , FITC , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (STAR77...)	HRP
Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Rabbit Anti Mouse IgG (STAR8...)	DyLight®800
Goat Anti Mouse IgG (STAR70...)	FITC
Rabbit Anti Mouse IgG (STAR13...)	HRP
Human Anti Mouse IgG1 (HCA036...)	HRP

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com "M283349:160301"	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

Printed on 01 May 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)