

Datasheet: MCA699PET

Description:	RAT ANTI HUMAN CD49f:RPE
Specificity:	CD49f
Other names:	INTEGRIN ALPHA 6 CHAIN, VLA-6
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	NKI-GoH3
Isotype:	IgG2a
Quantity:	25 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat - 1/5

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own systems using appropriate negative/positive controls.

Target Species	Human								
Species Cross Reactivity	Reacts with: Mouse, Dog, Pig, Cynomolgus monkey, Sheep N.B. Antibody reactivity and working conditions may vary between species.								
Product Form	Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized								
Reconstitution	Reconstitute in 0.25 ml distilled water								
Max Ex/Em	<table border="1"> <thead> <tr> <th>Fluorophore</th> <th>Excitation Max (nm)</th> <th>Emission Max (nm)</th> </tr> </thead> <tbody> <tr> <td>RPE 488nm laser</td> <td>496</td> <td>578</td> </tr> </tbody> </table>	Fluorophore	Excitation Max (nm)	Emission Max (nm)	RPE 488nm laser	496	578		
Fluorophore	Excitation Max (nm)	Emission Max (nm)							
RPE 488nm laser	496	578							
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant.								
Buffer Solution	Phosphate buffered saline								
Preservative	0.09% Sodium Azide								
Stabilisers	1% Bovine Serum Albumin 5% Sucrose								
Immunogen	BALB/c mouse mammary tumor cells								

**External Database
Links**

UniProt:

[P23229](#) [Related reagents](#)

Entrez Gene:

[3655](#) ITGA6 [Related reagents](#)

Fusion Partners

Spleen cells from immunized Sprague-Dawley rats were fused with cells of the SP2/0 mouse myeloma cell line

Specificity

Rat anti Human CD49f antibody, clone NKI-GoH3 recognizes CD49f, also known as the VLA-6 alpha chain. CD49f is a 1107 amino acid ~120 kDa cell surface glycoprotein that forms distinct complexes with CD29 (VLA beta-chain), resulting in the VLA-6 (alpha-6 beta-1) complex, expressed on human platelets, or with the beta-4 integrin resulting in the alpha-6 beta-4 complex expressed on various human epithelial cells.

Rat anti Human CD49f antibody, clone NKI-GoH3 reacts with platelets, megakaryocytes, T lymphocytes and common acute lymphoblastic leukemia cells (alpha-6 beta-1). In immunohistology the monoclonal antibody reacts with epithelial cells of a variety of tissues, peripheral nerves, microvascular endothelial cells, placenta cyto- and syncytiotrophoblasts. VLA-6 is an important mediator of cell binding to laminin.

Rat anti Human CD49f antibody, clone NKI-GoH3 blocks the binding of cells to the E8 fragment of laminin ([Sonnenberg et al. 1998](#)).

Flow Cytometry

Use 10ul of the suggested working dilution to label 10⁶ platelets in 100ul.

References

1. Jensen, K.B. *et al.* (2010) Assaying proliferation and differentiation capacity of stem cells using disaggregated adult mouse epidermis. [Nat Protoc. 5 \(5\): 898-911.](#)
2. Soligo, D. *et al.* (1989) Immunohistochemical reactivity on bone marrow and tissues of anti-VLA antibodies in the platelet panel, in Leucocyte Typing IV: White Cell Differentiation Antigens. Edited by Knapp, W. *et al.* Oxford University Press p1029-1032.
3. Sonnenberg, A. *et al.* (1986) Development of mouse mammary gland: identification of stages in differentiation of luminal and myoepithelial cells using monoclonal antibodies and polyvalent antiserum against keratin. [J Histochem Cytochem. 34 \(8\): 1037-46.](#)
4. Sonnenberg, A. *et al.* (1987) A complex of platelet glycoproteins Ic and IIa identified by a rat monoclonal antibody. [J Biol Chem. 262 \(21\): 10376-83.](#)
5. Hemler, M.E. *et al.* (1988) Multiple very late antigen (VLA) heterodimers on platelets. Evidence for distinct VLA-2, VLA-5 (fibronectin receptor), and VLA-6 structures. [J Biol Chem. 263 \(16\): 7660-5.](#)
6. Galkowska, H. *et al.* (1996) Reactivity of antibodies directed against human antigens with surface markers on canine leukocytes. [Vet Immunol Immunopathol. 53 \(3-4\): 329-34.](#)
7. Sonnenberg, A. *et al.* (1988) Laminin receptor on platelets is the integrin VLA-6. [Nature. 336 \(6198\): 487-9.](#)
8. Sonnenberg, A. *et al.* (1990) Integrin recognition of different cell-binding fragments of laminin (P1, E3, E8) and evidence that alpha 6 beta 1 but not alpha 6 beta 4 functions as a major receptor for fragment E8. [J Cell Biol. 110 \(6\): 2145-55.](#)
9. Yoshino, N. *et al.* (2000) Upgrading of flow cytometric analysis for absolute counts, cytokines and other antigenic molecules of cynomolgus monkeys (*Macaca fascicularis*) by using anti-human cross-reactive antibodies. [Exp Anim. 49 \(2\): 97-110.](#)
10. Sonnenberg, A. *et al.* (1990) The alpha 6 beta 1 (VLA-6) and alpha 6 beta 4 protein complexes: tissue distribution and biochemical properties. [J Cell Sci. 96 \(Pt 2\): 207-17.](#)
11. Sonnenberg, A. *et al.* (1988) Identification and characterization of a novel antigen complex on

mouse mammary tumor cells using a monoclonal antibody against platelet glycoprotein I α . [J Biol Chem. 263 \(28\): 14030-8.](#)

12. Le Bellego, F. *et al.* (2005) Cytoskeleton reorganization mediates α 6 β 1 integrin-associated actions of laminin on proliferation and survival, but not on steroidogenesis of ovine granulosa cells. [Reprod Biol Endocrinol. 3: 19.](#)

13. Anderson, C. *et al.* (2009) Sonic hedgehog-dependent synthesis of laminin α 1 controls basement membrane assembly in the myotome. [Development. 136: 3495-504.](#)

14. Collins, C.A. *et al.* (2011) Reprogramming adult dermis to a neonatal state through epidermal activation of β -catenin [Development. 138: 5189-99.](#)

15. Moreira, M. L. *et al.* (2016) Vaccination against canine leishmaniosis increases the phagocytic activity, nitric oxide production and expression of cell activation/migration molecules in neutrophils and monocytes. [Veterinary Parasitology. 15 Feb \[Epub ahead of print\]](#)

16. Mastrogianaki M *et al.* (2016) β -catenin stabilization in skin fibroblasts causes fibrotic lesions by preventing adipocyte differentiation of the reticular dermis. [J Invest Dermatol. pii: S0022-202X\(16\)00489-9. \[Epub ahead of print\]](#)

17. Schäfer, G. *et al.* (2013) The role of inflammation in HPV infection of the Oesophagus. [BMC Cancer. 13: 185.](#)

18. Peuhu, E. *et al.* (2017) Integrin beta 1 inhibition alleviates the chronic hyperproliferative dermatitis phenotype of SHARPIN-deficient mice [PLOS ONE. 12 \(10\): e0186628.](#)

19. Rayagiri, S.S. *et al.* (2018) Basal lamina remodeling at the skeletal muscle stem cell niche mediates stem cell self-renewal. [Nat Commun. 9 \(1\): 1075.](#)

Further Reading 1. Piriou-Guzylack, L. (2008) Membrane markers of the immune cells in swine: an update. [Vet Res. 39: 54.](#)

Storage Prior to reconstitution store at +4°C. Following reconstitution store at +4°C.

DO NOT FREEZE.

This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life 12 months from date of reconstitution.

Health And Safety Information Material Safety Datasheet documentation #10075 available at: 10075: <https://www.bio-rad-antibodies.com/uploads/MSDS/10075.pdf>

Regulatory For research purposes only

Related Products

Recommended Negative Controls

[RAT IgG2a NEGATIVE CONTROL:RPE \(MCA6005PE\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

Printed on 01 Aug 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)