

Datasheet: MCA691XZ

Description:	MOUSE IgG2b NEGATIVE CONTROL:Preservative Free
Specificity:	MOUSE IgG2b NEGATIVE CONTROL
Format:	Preservative Free
Product Type:	Negative/Isotype Control
Isotype:	IgG2b
Quantity:	1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			*
Immunohistology - Frozen	■			*
Immunohistology - Paraffin	■			*
Immunohistology - Resin	■			*
Radioimmunoassays	■			*

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only.

*It is recommended that the user adjusts the concentration of this reagent to be the same as the test reagents.

Target Species	Negative Control
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	None present
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml

Specificity

Mouse IgG2b Negative Control is negative on all human cells and cell lines tested. This antibody recognises a rat cell surface marker, and therefore cannot be used as a negative control in this species.

Test results show that MCA691C is also suitable for use as a negative control with bovine, ovine, porcine, canine and guinea-pig tissues.

Flow Cytometry

Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul.

References

1. Grant, A.J. *et al.* (2002) Hepatic expression of secondary lymphoid chemokine (CCL21) promotes the development of portal-associated lymphoid tissue in chronic inflammatory liver disease. [Am J Pathol. 160 \(4\): 1445-55.](#)
2. Zheng, X. *et al.* (2002) Interleukin-3, but not granulocyte-macrophage colony-stimulating factor and interleukin-5, inhibits apoptosis of human basophils through phosphatidylinositol 3-kinase: requirement of NF-KappaB-dependent and-independent pathways. [Immunology. 107:306-315.](#)
3. Dalli, J. *et al.* (2008) Annexin 1 mediates the rapid anti-inflammatory effects of neutrophil-derived microparticles. [Blood. 112 \(6\): 2512-9.](#)
4. Kapetanovic, R. *et al.* (2012) Pig bone marrow-derived macrophages resemble human macrophages in their response to bacterial lipopolysaccharide. [J Immunol. 188: 3382-94.](#)
5. Shoham, T. *et al.* (2001) Reduced expression of activin A in focal lymphoid agglomerates within nasal polyps. [J Histochem Cytochem. 49 \(10\): 1245-52.](#)

Storage

Store at -20°C only.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life

18 months from date of despatch.

Health And Safety Information

Material Safety Datasheet documentation #10162 available at:
10162: <https://www.bio-rad-antibodies.com/uploads/MSDS/10162.pdf>

Regulatory

For research purposes only

North & South America

Tel: +1 800 265 7376

Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M268243:150204'

Printed on 21 Jun 2018