

Datasheet: MCA749S

Description:	MOUSE ANTI GUINEA PIG CD4
Specificity:	CD4
Format:	Con S/N
Product Type:	Monoclonal Antibody
Clone:	CT7
Isotype:	IgG1
Quantity:	0.25 ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			1/10 - 1/100
Immunohistology - Frozen	■			1/10 - 1/100
Immunohistology - Paraffin			■	
ELISA			■	
Immunoprecipitation			■	
Western Blotting			■	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Guinea Pig
Product Form	Concentrated tissue Culture Supernatant - liquid
Preservative Stabilisers	0.1% Sodium Azide (NaN ₃) 0.7% Bovine Serum Albumin
Immunogen	Guinea pig peritoneal T-cells.
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the X63.Ag8.653 mouse myeloma cell line.
Specificity	Mouse anti Guinea Pig CD4 antibody, clone CT7 recognizes the CD4 antigen present on T Helper/Inducer lymphocytes.

References

1. Tan, B.T.G. *et al.* (1985) Production of monoclonal antibodies defining guinea pig T-cell surface markers and a strain 13 Ia-like antigen, the value of immunohistological screening. [Hybridoma 4: 115-124](#).
2. Baker, D. *et al.* (1987) Changes in lymphocyte subsets after treatment with cyclophosphamide

- and during the development of contact sensitivity in the guinea pig. [Int. J. Immunopharmac. 9: 175-183.](#)
3. Liversidge, J. *et al.* (1988) Experimental autoimmune uveitis (EAU): Immunophenotypic analysis of inflammatory cells in chorio retinal lesions: [Current eye research 7: 1231-1240.](#)
 4. Steerenburg, P.A. *et al.* (1991) Tumour rejection after transfer of line-10-immunity is mediated by two T-cell populations. [Cancer Immun. Immunother. 34: 103-110.](#)
 5. Debout, C. *et al.* (1991) The Kurloff Cell in estrogenized guinea pigs as a CT7+ 8BE6-CT6-MR-1-CT10-IgM-Lymphocyte with natural killer activity. [Natural Immunity and Cell Growth Regulation 10: 327-335.](#)
 6. Shang, S. *et al.* (2011) Activities of TMC207, rifampin, and pyrazinamide against Mycobacterium tuberculosis infection in guinea pigs. [Antimicrob Agents Chemother. 55 \(1\): 124-31.](#)
 7. Lacy, H.M. *et al.* (2011) Essential role for neutrophils in pathogenesis and adaptive immunity in *Chlamydia caviae* ocular infections. [Infect Immun. 79 \(5\): 1889-97.](#)
 8. Komori, T. *et al.* (2011) A Microbial Glycolipid Functions as a New Class of Target Antigen for Delayed-type Hypersensitivity. [J Biol Chem. 286: 16800-6.](#)
 9. Jeevan, A. *et al.* (2003) Differential expression of gamma interferon mRNA induced by attenuated and virulent Mycobacterium tuberculosis in guinea pig cells after Mycobacterium bovis BCG vaccination. [Infect Immun. 71: 354-64.](#)
 10. Schleiss, M.R. *et al.* (2007) Preconceptual administration of an alphavirus replicon UL83 (pp65 homolog) vaccine induces humoral and cellular immunity and improves pregnancy outcome in the guinea pig model of congenital cytomegalovirus infection. [J Infect Dis. 195: 789-98.](#)
 11. Turner, O.C. *et al.* (2003) Immunopathogenesis of pulmonary granulomas in the guinea pig after infection with Mycobacterium tuberculosis. [Infect Immun. 71: 864-71.](#)
 12. Wang, Y. *et al.* (2010) Local host response to chlamydial urethral infection in male guinea pigs. [Infect Immun. 78: 1670-81.](#)
 13. Mishra, N.C. *et al.* (2010) Sulfur mustard induces immune sensitization in hairless guinea pigs. [Int Immunopharmacol. 10: 193-9.](#)
 14. Hiromatsu, K. *et al.* (2002) Induction of CD1-restricted immune responses in guinea pigs by immunization with mycobacterial lipid antigens. [J Immunol. 169: 330-9.](#)
 15. Dascher, C.C. *et al.* (1999) Conservation of a CD1 multigene family in the guinea pig. [J Immunol. 163: 5478-88.](#)
 16. Rousseau, C. *et al.* (2003) Sulfolipid Deficiency Does Not Affect the Virulence of Mycobacterium tuberculosis H37Rv in Mice and Guinea Pigs [Infect Immun. 71: 4684-90.](#)
 17. Kramp, J.C. *et al.* (2011) The in vivo immunomodulatory effect of recombinant tumour necrosis factor-alpha in guinea pigs vaccinated with Mycobacterium bovis bacille Calmette-Guérin. [Clin Exp Immunol. 165: 110-20.](#)
 18. Chitano, P. *et al.* (2014) Ovalbumin sensitization of guinea pig at birth prevents the ontogenetic decrease in airway smooth muscle responsiveness. [Physiol Rep. 2 \(12\): .](#)
 19. Gupta, A. *et al.* (2012) Efficacy of Mycobacterium indicus pranii immunotherapy as an adjunct to chemotherapy for tuberculosis and underlying immune responses in the lung. [PLoS One. 7 \(7\): e39215.](#)
 20. Podell, B.K. *et al.* (2014) Increased severity of tuberculosis in Guinea pigs with type 2 diabetes: a model of diabetes-tuberculosis comorbidity. [Am J Pathol. 184 \(4\): 1104-18.](#)
 21. Shang, S. *et al.* (2012) Drug treatment combined with BCG vaccination reduces disease reactivation in guinea pigs infected with Mycobacterium tuberculosis. [Vaccine. 30 \(9\): 1572-82.](#)
 22. Yang H *et al.* (2011) Three protein cocktails mediate delayed-type hypersensitivity responses indistinguishable from that elicited by purified protein derivative in the guinea pig model of Mycobacterium tuberculosis infection. [Infect Immun. 79 \(2\): 716-23.](#)
 23. Jeevan A *et al.* (2013) Guinea pig skin, a model for epidermal cellular and molecular changes induced by UVR *in vivo* and *in vitro*: effects on Mycobacterium bovis Bacillus Calmette-Guérin vaccination. [Photochem Photobiol. 89 \(1\): 189-98.](#)
 24. Miszczyk, E. *et al.* (2014) Antigen-specific lymphocyte proliferation as a marker of immune response in guinea pigs with sustained Helicobacter pylori infection. [Acta Biochim Pol. 61 \(2\):](#)

Storage

Store at +4°C for one month or at -20°C for longer.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life

18 months from date of despatch.

Health And Safety Information

Material Safety Datasheet documentation #10495 available at:
10495: <https://www.bio-rad-antibodies.com/uploads/MSDS/10495.pdf>

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87...)	Alk. Phos. , HRP
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®549 , DyLight®649 , DyLight®680 , DyLight®800 , FITC , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (STAR77...)	HRP
Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Rabbit Anti Mouse IgG (STAR8...)	DyLight®800
Goat Anti Mouse IgG (STAR70...)	FITC
Rabbit Anti Mouse IgG (STAR13...)	HRP
Human Anti Mouse IgG1 (HCA036...)	HRP

North & South America Tel: +1 800 265 7376
Fax: +1 919 878 3751
Email: antibody_sales_us@bio-rad.com

Worldwide Tel: +44 (0)1865 852 700
Fax: +44 (0)1865 852 739
Email: antibody_sales_uk@bio-rad.com

Europe Tel: +49 (0) 89 8090 95 21
Fax: +49 (0) 89 8090 95 50
Email: antibody_sales_de@bio-rad.com

'M304117:170317'

Printed on 01 May 2018