

Datasheet: MCA740B

Description:	MOUSE ANTI HUMAN CD42b:Biotin
Specificity:	CD42b
Other names:	GPIB-ALPHA
Format:	Biotin
Product Type:	Monoclonal Antibody
Clone:	AK2
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. The suggested working dilution is given as a guide only. It is recommended that the user titrates the antibody for use in his/her own system using appropriate negative/positive controls.

Target Species	Human
Product Form	Purified IgG conjugated to Biotin - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant.
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide 1% Bovine Serum Albumin
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml
External Database Links	UniProt: P07359 Related reagents Entrez Gene: 2811 GP1BA Related reagents
Specificity	Mouse anti Human CD42b antibody, clone AK2 recognizes the human CD42b cell surface antigen, also known as platelet glycoprotein GP1B. CD42b is expressed by platelets and

megakaryocytes.

Clone AK2 has been reported to block the binding of von Willebrand Factor (VWF) to platelets

Flow Cytometry

Use 10ul of the suggested working dilution to label 100ul whole blood.

References

1. Ward, C.M. & Berndt, M.C. (1995) Epitope and functional characterization of the CD42 (gplb/IX) mAb panel. *Leucocyte Typing V. White Cell Differentiation Antigens. Volume Two.* Oxford University Press, Oxford.
2. Burgess, J.K. *et al.* (1998) Quinine-dependent antibodies bind a restricted set of epitopes on the glycoprotein Ib-IX complex: characterization of the epitopes. [Blood. 92: 2366-73.](#)
3. Burgess, J.K. *et al.* (2000) Rifampicin-dependent antibodies bind a similar or identical epitope to glycoprotein IX-specific quinine-dependent antibodies. [Blood. 95: 1988-92.](#)
4. Jayo, A. *et al.* (2010) L718P mutation in the membrane-proximal cytoplasmic tail of beta 3 promotes abnormal alpha IIb beta 3 clustering and lipid microdomain coalescence, and associates with a thrombasthenia-like phenotype. [Haematologica. 95: 1158-66.](#)
5. Lova, P. *et al.* (2004) Contribution of protease-activated receptors 1 and 4 and glycoprotein Ib-IX-V in the G(i)-independent activation of platelet Rap1B by thrombin. [J Biol Chem. 279: 25299-306.](#)
6. Shen, Y. *et al.* (2000) Requirement of leucine-rich repeats of glycoprotein (GP) Ibalpha for shear-dependent and static binding of von Willebrand factor to the platelet membrane GP Ib-IX-V complex. [Blood. 95: 903-10.](#)
7. Tait, A.S. *et al.* (2002) Site-directed mutagenesis of platelet glycoprotein Ib alpha demonstrating residues involved in the sulfation of tyrosines 276, 278, and 279. [Blood. 99: 4422-7.](#)
8. Yap, C.L. *et al.* (2000) Synergistic adhesive interactions and signaling mechanisms operating between platelet glycoprotein Ib/IX and integrin alpha IIb beta 3. Studies in human platelets and transfected Chinese hamster ovary cells. [J Biol Chem. 275: 41377-88.](#)
9. Wright, S.D. *et al.* (1993) Double heterozygosity for mutations in the platelet glycoprotein IX gene in three siblings with Bernard-Soulier syndrome. [Blood. 81: 2339-47.](#)
10. Nomura, S. *et al.* (1995) Significance of cytokines and CD68-positive microparticles in immune thrombocytopenic purpura. [Eur J Haematol. 55: 49-56.](#)
11. Speich, H.E. *et al.* (2008) Platelets undergo phosphorylation of Syk at Y525/526 and Y352 in response to pathophysiological shear stress. [Am J Physiol Cell Physiol. 295: C1045-54.](#)
12. Amor, N.B. *et al.* (2009) Acidic-store depletion is required for human platelet aggregation. [Blood Coagul Fibrinolysis. 20: 511-6.](#)
13. Tasneem, S. *et al.* (2009) Platelet adhesion to multimerin 1 in vitro: influences of platelet membrane receptors, von Willebrand factor and shear. [J Thromb Haemost. 7: 685-92.](#)
14. Lincoln, B. *et al.* (2010) Integrated system investigating shear-mediated platelet interactions with von Willebrand factor using microliters of whole blood [Anal Biochem. 405: 174-83.](#)
15. Balduini, A. *et al.* (2008) Adhesive receptors, extracellular proteins and myosin IIA orchestrate proplatelet formation by human megakaryocytes. [J Thromb Haemost. 6: 1900-7.](#)
16. Goetzl EJ *et al.* (2016) Human plasma platelet-derived exosomes: effects of aspirin. [FASEB J. Feb 12. pii: fj.201500150R. \[Epub ahead of print\]](#)
17. Michalska-Jakubus, M. *et al.* (2016) Plasma endothelial microparticles reflect the extent of capillaroscopic alterations and correlate with the severity of skin involvement in systemic sclerosis. [Microvasc Res. Nov 23. pii: S0026-2862\(16\)30097-8. \[Epub ahead of print\]](#)
18. Ralph, A. *et al.* (2016) Computational Tracking of Shear-Mediated Platelet Interactions with von Willebrand Factor. [Cardiovasc Eng Technol. 7 \(4\): 389-405.](#)
19. Rossi, E. *et al.* (2017) Human endoglin as a potential new partner involved in platelet-endothelium interactions. [Cell Mol Life Sci. Oct 28 \[Epub ahead of print\].](#)

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life	18 months from date of despatch.
-------------------	----------------------------------

Health And Safety Information	Material Safety Datasheet documentation #10041 available at: 10041: https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf
--------------------------------------	--

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

'M304753:170412'

Printed on 05 May 2018