

Datasheet: MCA740GA

Description:	MOUSE ANTI HUMAN CD42b
Specificity:	CD42b
Other names:	GPIB-ALPHA
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	AK2
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			1/50 - 1/100
ELISA	■			
Immunoprecipitation	■			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. The suggested working dilution is given as a guide only. It is recommended that the user titrates the antibody for use in his/her own system using appropriate negative/positive controls.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1 mg/ml
External Database Links	UniProt: P07359 Related reagents Entrez Gene:

Specificity	Mouse anti Human CD42b antibody, clone AK2 recognizes the human CD42b cell surface antigen, also known as platelet glycoprotein GP1B. CD42b is expressed by platelets and megakaryocytes. Clone AK2 has been reported to block the binding of von Willebrand Factor (VWF) to platelets
Flow Cytometry	Use 10ul of the suggested working dilution to label 100ul whole blood.
References	1. Ward, C.M. & Berndt, M.C. (1995) Epitope and functional characterization of the CD42 (gplb/IX) mAb panel. Leucocyte Typing V. White Cell Differentiation Antigens. Volume Two. Oxford University Press, Oxford. 2. Burgess, J.K. <i>et al.</i> (1998) Quinine-dependent antibodies bind a restricted set of epitopes on the glycoprotein Ib-IX complex: characterization of the epitopes. Blood. 92: 2366-73. 3. Burgess, J.K. <i>et al.</i> (2000) Rifampicin-dependent antibodies bind a similar or identical epitope to glycoprotein IX-specific quinine-dependent antibodies. Blood. 95: 1988-92. 4. Jayo, A. <i>et al.</i> (2010) L718P mutation in the membrane-proximal cytoplasmic tail of beta 3 promotes abnormal alpha IIb beta 3 clustering and lipid microdomain coalescence, and associates with a thrombasthenia-like phenotype. Haematologica. 95: 1158-66. 5. Lova, P. <i>et al.</i> (2004) Contribution of protease-activated receptors 1 and 4 and glycoprotein Ib-IX-V in the G(i)-independent activation of platelet Rap1B by thrombin. J Biol Chem. 279: 25299-306. 6. Shen, Y. <i>et al.</i> (2000) Requirement of leucine-rich repeats of glycoprotein (GP) Ibalpha for shear-dependent and static binding of von Willebrand factor to the platelet membrane GP Ib-IX-V complex. Blood. 95: 903-10. 7. Tait, A.S. <i>et al.</i> (2002) Site-directed mutagenesis of platelet glycoprotein Ib alpha demonstrating residues involved in the sulfation of tyrosines 276, 278, and 279. Blood. 99: 4422-7. 8. Yap, C.L. <i>et al.</i> (2000) Synergistic adhesive interactions and signaling mechanisms operating between platelet glycoprotein Ib/IX and integrin alpha IIbbeta 3. Studies in human platelets and transfected Chinese hamster ovary cells. J Biol Chem. 275: 41377-88. 9. Wright, S.D. <i>et al.</i> (1993) Double heterozygosity for mutations in the platelet glycoprotein IX gene in three siblings with Bernard-Soulier syndrome. Blood. 81: 2339-47. 10. Nomura, S. <i>et al.</i> (1995) Significance of cytokines and CD68-positive microparticles in immune thrombocytopenic purpura. Eur J Haematol. 55: 49-56. 11. Speich, H.E. <i>et al.</i> (2008) Platelets undergo phosphorylation of Syk at Y525/526 and Y352 in response to pathophysiological shear stress. Am J Physiol Cell Physiol. 295: C1045-54. 12. Amor, N.B. <i>et al.</i> (2009) Acidic-store depletion is required for human platelet aggregation. Blood Coagul Fibrinolysis. 20: 511-6. 13. Tasneem, S. <i>et al.</i> (2009) Platelet adhesion to multimerin 1 in vitro: influences of platelet membrane receptors, von Willebrand factor and shear. J Thromb Haemost. 7: 685-92. 14. Lincoln, B. <i>et al.</i> (2010) Integrated system investigating shear-mediated platelet interactions with von Willebrand factor using microliters of whole blood Anal Biochem. 405: 174-83. 15. Balduini, A. <i>et al.</i> (2008) Adhesive receptors, extracellular proteins and myosin IIA orchestrate proplatelet formation by human megakaryocytes. J Thromb Haemost. 6: 1900-7. 16. Goetzl EJ <i>et al.</i> (2016) Human plasma platelet-derived exosomes: effects of aspirin. FASEB J. Feb 12. pii: fj.201500150R. [Epub ahead of print] 17. Michalska-Jakubus, M. <i>et al.</i> (2016) Plasma endothelial microparticles reflect the extent of capillaroscopic alterations and correlate with the severity of skin involvement in systemic sclerosis. Microvasc Res. Nov 23. pii: S0026-2862(16)30097-8. [Epub ahead of print] 18. Ralph, A. <i>et al.</i> (2016) Computational Tracking of Shear-Mediated Platelet Interactions with von Willebrand Factor. Cardiovasc Eng Technol. 7 (4): 389-405.

19. Rossi, E. *et al.* (2017) Human endoglin as a potential new partner involved in platelet-endothelium interactions. [Cell Mol Life Sci. Oct 28 \[Epub ahead of print\]](#).

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life

18 months from date of despatch.

Health And Safety Information

Material Safety Datasheet documentation #10040 available at:
10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87...)	Alk. Phos. , HRP
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®549 , DyLight®649 , DyLight®680 , DyLight®800 , FITC , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (STAR77...)	HRP
Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Rabbit Anti Mouse IgG (STAR8...)	DyLight®800
Goat Anti Mouse IgG (STAR70...)	FITC
Rabbit Anti Mouse IgG (STAR13...)	HRP
Human Anti Mouse IgG1 (HCA036...)	HRP

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South America Tel: +1 800 265 7376
Fax: +1 919 878 3751
Email: antibody_sales_us@bio-rad.com

Worldwide Tel: +44 (0)1865 852 700
Fax: +44 (0)1865 852 739
Email: antibody_sales_uk@bio-rad.com

Europe Tel: +49 (0) 89 8090 95 21
Fax: +49 (0) 89 8090 95 50
Email: antibody_sales_de@bio-rad.com

'M316290:180503'

Printed on 05 May 2018