

Datasheet: MCA743EL

Description:	MOUSE ANTI HUMAN CD49b:Low Endotoxin
Specificity:	CD49b
Other names:	INTEGRIN ALPHA 2 CHAIN, VLA-2
Format:	Low Endotoxin
Product Type:	Monoclonal Antibody
Clone:	AK7
Isotype:	IgG1
Quantity:	0.5 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			1/50 - 1/100
Immunohistology - Frozen (1)	■			1/500 - 1/1000
Immunohistology - Paraffin		■		
ELISA			■	
Immunoprecipitation			■	
Western Blotting			■	
Functional Assays	■			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1)The epitope recognised by this antibody is reported to be sensitive to formaldehyde fixation and tissue processing. Bio-Rad recommends the use of acetone fixation for frozen sections.

Target Species	Human
Species Cross Reactivity	Reacts with: Cynomolgus monkey, Rhesus Monkey, Baboon N.B. Antibody reactivity and working conditions may vary between species.
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	None present

Carrier Free	Yes
Endotoxin Level	<0.01EU/ug
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
External Database Links	UniProt: P17301 Related reagents Entrez Gene: 3673 ITGA2 Related reagents
Synonyms	CD49B
Fusion Partners	Spleen cells from immunised mice were fused with cells of the mouse NS1 myeloma cell line.
Specificity	Mouse anti Human CD49b antibody, clone AK7 recognizes the integrin alpha 2 subunit, a ~160 kDa glycoprotein that non-covalently associates with the ~130 kDa integrin beta 1 subunit to form the VLA-2 complex. CD49b is expressed by platelets, long term cultivated T cells, approximately 50% of monocytes and most adherent cell lines. Mouse anti Human CD49b antibody, clone AK7 inhibits cell attachment to collagen.
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	1. Favaloro, E.J. <i>et al.</i> (1990) Co-expression of haemopoietic antigens on vascular endothelial cells: a detailed phenotypic analysis. Br J Haematol. 74 (4): 385-94. 2. Mazurov, A.V. <i>et al.</i> (1991) A monoclonal antibody, VM64, reacts with a 130 kDa glycoprotein common to platelets and endothelial cells: heterogeneity in antibody binding to human aortic endothelial cells. Thromb Haemost. 66 (4): 494-9. 3. Yoshino N <i>et al.</i> (2000) Upgrading of flow cytometric analysis for absolute counts, cytokines and other antigenic molecules of cynomolgus monkeys (<i>Macaca fascicularis</i>) by using anti-human cross-reactive antibodies. Exp Anim. 49 (2): 97-110. 4. Ciarlet, M. <i>et al.</i> (2002) VLA-2 (alpha2beta1) integrin promotes rotavirus entry into cells but is not necessary for rotavirus attachment. J Virol. 76 (3): 1109-23. 5. Cavers, M. <i>et al.</i> (2002) Differential expression of beta1 and beta2 integrins and L-selectin on CD4+ and CD8+ T lymphocytes in human blood: comparative analysis between isolated cells, whole blood samples and cryopreserved preparations. Clin Exp Immunol. 127: 60-5. 6. Mörtberg A <i>et al.</i> (2016) Sensitive detection of platelet-specific antibodies with a modified MAIPA using biotinylated antibodies and streptavidin-coated beads. J Immunol Methods. 434: 9-15. 7. Rane, J.K. <i>et al.</i> (2016) Inhibition of the glucocorticoid receptor results in an enhanced miR-99a/100-mediated radiation response in stem-like cells from human prostate cancers. Oncotarget. 7 (32): 51965-80. 8. Koutsogiannaki S. & Kaloyianni M. (2011) Effect of 17β-estradiol on adhesion of <i>Mytilus galloprovincialis</i> hemocytes to selected substrates. Role of alpha2 integrin subunit Fish & Shellfish Immunology. 31 (1): 73-80. 9. Hamaia, S.W. <i>et al.</i> (2012) Mapping of potent and specific binding motifs, GLOGEN and GVOGEA, for integrin α1β1 using collagen toolkits II and III. J Biol Chem. 287 (31): 26019-28. 10. Koutsogiannaki, S. & Kaloyianni, M. (2011) Effect of 17β-estradiol on adhesion of <i>Mytilus</i>

galloprovincialis hemocytes to selected substrates. Role of alpha2 integrin subunit. [Fish Shellfish Immunol. 31 \(1\): 73-80.](#)

Storage

Store at -20°C only.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life

18 months from date of despatch.

Health And Safety Information

Material Safety Datasheet documentation #10162 available at:
10162: <https://www.bio-rad-antibodies.com/uploads/MSDS/10162.pdf>

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87...)	Alk. Phos. , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (STAR77...)	HRP
Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Rabbit Anti Mouse IgG (STAR8...)	DyLight®800
Goat Anti Mouse IgG (STAR70...)	FITC
Rabbit Anti Mouse IgG (STAR13...)	HRP
Human Anti Mouse IgG1 (HCA036...)	HRP
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®549 , DyLight®649 , DyLight®680 , DyLight®800 , FITC , HRP

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:Low Endotoxin \(MCA928EL\)](#)

North & South America Tel: +1 800 265 7376
Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700
Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21
Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M318177:180718'

Printed on 01 Aug 2018