

Datasheet: MCA775

Description:	MOUSE ANTI RAT CD18
Specificity:	CD18
Other names:	INTEGRIN BETA 2 CHAIN
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	WT.3
Isotype:	IgG1
Quantity:	0.2 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			1/10 - 1/50
Immunohistology - Frozen	■			
Immunohistology - Paraffin			■	
ELISA			■	
Immunoprecipitation	■			
Western Blotting			■	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Rat
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1 mg/ml
Immunogen	FTL-43.

Fusion Partners	Lymph node cells from an immunized BALB/c mouse were fused with cells of the PAI mouse myeloma cell line.
Specificity	Mouse anti Rat CD18 antibody, clone WT.3 reacts with the rat beta 2 integrin, which is designated CD18. The beta 2 integrin may be associated with the integrin alphaL, alpha^X or alphaM subunits. CD18 is expressed by all leucocytes. Mouse anti Rat CD18 antibody has been shown to inhibit homotypic aggregation of PHA blasts. It also blocks binding of rat lymphocytes to purified rat ICAM-1.
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	1. Tamatani, T. & Miyasaka, M. (1990) Identification of monoclonal antibodies reactive with the rat homolog of ICAM-1, and evidence for a differential involvement of ICAM-1 in the adherence of resting versus activated lymphocytes to high endothelial cells. Int Immunol. 2 (2): 165-71. 2. Tamatani, T. <i>et al.</i> (1991) Characterization of the rat leukocyte integrin, CD11/CD18, by the use of LFA-1 subunit-specific monoclonal antibodies. Eur J Immunol. 21 (3): 627-33. 3. Tamatani, T. <i>et al.</i> (1991) Molecular mechanisms underlying lymphocyte recirculation. II. Differential regulation of LFA-1 in the interaction between lymphocytes and high endothelial cells. Eur J Immunol. 21 (3): 855-8. 4. Alvarez, A. <i>et al.</i> (2007) Gastric antisecretory drugs induce leukocyte-endothelial cell interactions through gastrin release and activation of CCK-2 receptors. J Pharmacol Exp Ther. 323: 406-413. 5. Wu JC <i>et al.</i> (1996) The relationship of adhesion molecules and leukocyte infiltration in chronic tubulointerstitial nephritis induced by puromycin aminonucleoside in Wistar rats. Clin Immunol Immunopathol. 79 (3): 229-35. 6. Nicholls, S.M. <i>et al.</i> (2006) Differences in leukocyte phenotype and interferon-gamma expression in stroma and endothelium during corneal graft rejection. Exp Eye Res. 83 (2): 339-47. 7. Ishida, S. <i>et al.</i> (2003) Leukocytes mediate retinal vascular remodeling during development and vaso-obliteration in disease. Nat Med. 9 (6): 781-8. 8. Fabian, R.H. & Kent, T.A. (1999) Superoxide anion production during reperfusion is reduced by an antineutrophil antibody after prolonged cerebral ischemia. Free Radic Biol Med. 26 (3-4): 355-61. 9. Shen K <i>et al.</i> (1995) Circulating leukocyte counts, activation, and degranulation in Dahl hypertensive rats. Circ Res. 76 (2): 276-83. 10. Natile-McMenemy, N. <i>et al.</i> (2007) Minocycline decreases in vitro microglial motility, beta1-integrin, and Kv1.3 channel expression. J Neurochem. 103 (5): 2035-46. 11. Martinelli, R. <i>et al.</i> (2009) ICAM-1-mediated endothelial nitric oxide synthase activation via calcium and AMP-activated protein kinase is required for transendothelial lymphocyte migration. Mol Biol Cell. 20 (3): 995-1005. 12. Herrmann, I.K. <i>et al.</i> (2015) Differentiating sepsis from non-infectious systemic inflammation based on microvesicle-bacteria aggregation. Nanoscale. 7 (32): 13511-20.
Storage	Store at +4°C or at -20°C if preferred. This product should be stored undiluted. Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Shelf Life	18 months from date of despatch.

**Health And Safety
Information**

Material Safety Datasheet documentation #10040 available at:
10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87...)	Alk. Phos. , HRP
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®549 , DyLight®649 , DyLight®680 , DyLight®800 , FITC , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (STAR77...)	HRP
Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Rabbit Anti Mouse IgG (STAR8...)	DyLight®800
Goat Anti Mouse IgG (STAR70...)	FITC
Rabbit Anti Mouse IgG (STAR13...)	HRP
Human Anti Mouse IgG1 (HCA036...)	HRP

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA1209\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
--------------------------------------	---	------------------	---	---------------	---

'M316311:180503'

Printed on 05 May 2018