

Datasheet: MCA773FB

Description:	MOUSE ANTI RAT CD54:FITC
Specificity:	CD54
Other names:	ICAM-1
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	1A29
Isotype:	IgG1
Quantity:	0.5 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			1/25 - 1/100

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Rat		
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide		
Stabilisers	1% Bovine Serum Albumin		
Approx. Protein Concentrations	IgG concentration 0.5 mg/ml		
Immunogen	Rat Ax cells (a HEV derived cell line).		
External Database Links	UniProt: Q00238 Related reagents		

Entrez Gene:[25464](#) Icam1 [Related reagents](#)

Synonyms	Icam-1
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells from the PAI mouse myeloma cell line.
Specificity	Mouse anti Rat CD54 antibody, clone 1A29 recognizes the rat CD54 cell surface antigen, also known as intercellular adhesion molecule-1 (ICAM-1), a ~90 kDa adhesion molecule belonging to the immunoglobulin superfamily. CD54 is a cell surface ligand of the lymphocyte integrin, LFA-1 and is known to play an important role in various cell-cell interactions in the immune system. Studies suggest that cross-linking of ICAM-1 using clone 1A29 induces calcium signalling (Etienne et al.). Functionally Mouse anti Rat CD54 antibody, clone 1A29 inhibits homotypic aggregation of PHA blasts.
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	1. Kawai, T. <i>et al.</i> (1999) Selective diapedesis of Th1 cells induced by endothelial cell RANTES. J Immunol. 163: 3269-78. 2. Sato, N. <i>et al.</i> (2000) Roles of ICAM-1 for abnormal leukocyte recruitment in the microcirculation of bleomycin-induced fibrotic lung injury. Am J Respir Crit Care Med. 161: 1681-8. 3. Etienne, S. <i>et al.</i> (1998) ICAM-1 signaling pathways associated with Rho activation in microvascular brain endothelial cells. J Immunol. 161 (10): 5755-61. 4. Etienne-Manneville, S. <i>et al.</i> (2000) ICAM-1-coupled cytoskeletal rearrangements and transendothelial lymphocyte migration involve intracellular calcium signaling in brain endothelial cell lines. J Immunol. 165 (6): 3375-83. 5. McKechnie, N. M. <i>et al.</i> (2002) Antigenic mimicry: Onchocerca volvulus antigen-specific T cells and ocular inflammation. Invest Ophthalmol Vis Sci. 43:411-8. 6. Banerjee, S. <i>et al.</i> (2003) Development of organised conjunctival leucocyte aggregates after corneal transplantation in rats. Br J Ophthalmol. 2003 Dec;87(12):1515-22. 7. Kielian, T. <i>et al.</i> (2000) Proinflammatory cytokine, chemokine, and cellular adhesion molecule expression during the acute phase of experimental brain abscess development. Am J Pathol. 157: 647-58. 8. Trinh, L. <i>et al.</i> (2008) The corneal endothelium in an endotoxin-induced uveitis model: correlation between <i>in vivo</i> confocal microscopy and immunohistochemistry. Mol Vis. 14: 1149-56. 9. Adamson, P. <i>et al.</i> (1999) Lymphocyte migration through brain endothelial cell monolayers involves signaling through endothelial ICAM-1 via a rho-dependent pathway. J Immunol. 162: 2964-73. 10. Arsenović-Ranin, N. <i>et al.</i> (2000) A monoclonal antibody to the rat Crry/p65 antigen, a complement regulatory membrane protein, stimulates adhesion and proliferation of thymocytes. Immunology. 100: 334-44. 11. Beck-Schimmer, B. <i>et al.</i> (2001) Hypoxia mediates increased neutrophil and macrophage adhesiveness to alveolar epithelial cells. Am J Respir Cell Mol Biol. 25: 780-7. 12. Couty, J.P. (2007) PECAM-1 engagement counteracts ICAM-1-induced signaling in brain vascular endothelial cells. J Neurochem. 103: 793-801. 13. Deng, H. <i>et al.</i> (2003) Mild hypothermia inhibits inflammation after experimental stroke and brain inflammation. Stroke. 34: 2495-501. 14. Ikezumi, Y. <i>et al.</i> (2004) Macrophage-mediated renal injury is dependent on signaling via the

- JNK pathway. [J Am Soc Nephrol. 15: 1775-84.](#)
15. Kanellis, J. *et al.* (2010) JNK signalling in human and experimental renal ischaemia/reperfusion injury. [Nephrol Dial Transplant. 25: 2898-908.](#)
 16. Westermann, D. *et al.* (2007) Cardioprotective and anti-inflammatory effects of interleukin converting enzyme inhibition in experimental diabetic cardiomyopathy. [Diabetes. 56: 1834-41.](#)
 17. Zhu, X. *et al.* (2003) Matrine protects sinusoidal endothelial cells from cold ischemia and reperfusion injury in rat orthotopic liver transplantation. [Ann Clin Lab Sci. 33: 216-25.](#)
 18. Choi, J.S. *et al.* (2011) Mild Hypothermia Attenuates Intercellular Adhesion Molecule-1 Induction via Activation of Extracellular Signal-Regulated Kinase-1/2 in a Focal Cerebral Ischemia Model. [Stroke Res Treat. 2011: 846716.](#)
 19. Azcutia V *et al.* (2010) Inflammation determines the pro-adhesive properties of high extracellular d-glucose in human endothelial cells *in vitro* and rat microvessels *in vivo*. [PLoS One. 5 \(4\): e10091.](#)
 20. Li, W. & Klein, S.L. (2012) Seoul virus-infected rat lung endothelial cells and alveolar macrophages differ in their ability to support virus replication and induce regulatory T cell phenotypes. [J Virol. 86 \(21\): 11845-55.](#)
 21. Gates, D. *et al.* (2012) Apo J/clusterin expression and secretion: evidence for 15-deoxy- $\Delta(12,14)$ -PGJ(2)-dependent mechanism. [Biochim Biophys Acta. 1821 \(2\): 335-42.](#)
 22. Li, Z. *et al.* (2015) Three-dimensional graphene foams loaded with bone marrow derived mesenchymal stem cells promote skin wound healing with reduced scarring. [Mater Sci Eng C Mater Biol Appl. 57: 181-8.](#)
 23. Gautier, S. *et al.* (2015) PPAR-Alpha Agonist Used at the Acute Phase of Experimental Ischemic Stroke Reduces Occurrence of Thrombolysis-Induced Hemorrhage in Rats. [PPAR Research. 2015: 1-6.](#)
 24. Liu, Y.C. *et al.* (2013) A biodegradable, sustained-released, prednisolone acetate microfilm drug delivery system effectively prolongs corneal allograft survival in the rat keratoplasty model. [PLoS One. 8 \(8\): e70419.](#)
 25. Ichihara, Y. *et al.* (2018) Self-assembling peptide hydrogel enables instant epicardial coating of the heart with mesenchymal stromal cells for the treatment of heart failure. [Biomaterials. 154: 12-23.](#)

Storage	Store at +4°C or at -20°C if preferred.
	This product should be stored undiluted.
	Storage in frost free freezers is not recommended. This product is photosensitive and should be protected from light.
	Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life	18 months from date of despatch.
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: 10041: https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA1209F\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com "M301700:170109"	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

Printed on 05 May 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)