

Datasheet: MCA796XZ

Description:	MOUSE ANTI HUMAN CD62P:Preservative Free
Specificity:	CD62P
Other names:	P-SELECTIN
Format:	Preservative Free
Product Type:	Monoclonal Antibody
Clone:	AK-6
Isotype:	IgG1
Quantity:	1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			1/10 - 1/100
Immunohistology - Frozen	■			1/1000 - 1/5000
Immunohistology - Paraffin (1)	■			1/500 - 1/200
ELISA			■	
Immunoprecipitation			■	
Western Blotting	■			
Immunofluorescence	■			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1) **Staining may be enhanced using Sodium Citrate buffer pH6.0**

Target Species	Human
Species Cross Reactivity	Reacts with: Rhesus Monkey N.B. Antibody reactivity and working conditions may vary between species.
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	None present
Carrier Free	Yes

Approx. Protein Concentrations	IgG concentration 1 mg/ml
Immunogen	Human platelet membrane glycoproteins.
External Database Links	UniProt: P16109 Related reagents Entrez Gene: 6403 SELP Related reagents
Synonyms	GMRP, GRMP
Specificity	Mouse anti Human CD62P antibody, clone AK-6 recognizes the CD62P, also known as P-selectin, Granule membrane protein 140, GMP140, Leukocyte-endothelial cell adhesion molecule 3 or Platelet activation dependent granule-external membrane protein. CD62P is a 830 amino acid, including a 41 amino acid signal peptide, ~140 kDa single pass type I transmembrane glycoprotein expressed on activated platelets and endothelial cell CD62P plays an important role in adhesive processes between leucocytes and endothelial cells. CD62P is a component of the platelet alpha granule and is rapidly translocated to the plasma membrane upon activation (Stenberg et al. 1985).
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
Histology Positive Control Tissue	Human Tonsil
References	- Dunlop, L.C. <i>et al.</i> (1992) Characterization of GMP-140 (P-selectin) as a circulating plasma protein. J Exp Med. 175 (4): 1147-50. - Skinner, M.P. <i>et al.</i> (1989) Characterization of human platelet GMP-140 as a heparin-binding protein. Biochem Biophys Res Commun. 164 (3): 1373-9. - Skinner, M.P. <i>et al.</i> (1991) GMP-140 binding to neutrophils is inhibited by sulfated glycans. J Biol Chem. 266 (9): 5371-4. - Sopper, S. <i>et al.</i> (1997) Lymphocyte subsets and expression of differentiation markers in blood and lymphoid organs of rhesus monkeys. Cytometry. 29 (4): 351-62. - Bevilacqua, M.P. & Nelson, R.M. (1993) Selectins. J Clin Invest. 91 (2): 379-87. - Roos-Engstrand, E. <i>et al.</i> (2005) Increased expression of p38 MAPK in human bronchial epithelium after lipopolysaccharide exposure. Eur Respir J. 25 (5): 797-803. - Kornerup, K.N. <i>et al.</i> (2010) Circulating platelet-neutrophil complexes are important for subsequent neutrophil activation and migration. J Appl Physiol. 109: 758-67. - Norling, L.V. <i>et al.</i> (2008) Inhibitory control of endothelial galectin-1 on in vitro and in vivo lymphocyte trafficking. FASEB J. 22: 682-90. - Dalli, J. <i>et al.</i> (2008) Annexin 1 mediates the rapid anti-inflammatory effects of neutrophil-derived microparticles. Blood. 112 (6): 2512-9. - Wassmer, S.C. <i>et al.</i> (2008) Platelet-induced clumping of Plasmodium falciparum-infected erythrocytes from Malawian patients with cerebral malaria-possible modulation in vivo by thrombocytopenia. J Infect Dis. 197: 72-8. - Theoret, J.F. <i>et al.</i> (2001) P-selectin antagonism with recombinant p-selectin glycoprotein ligand-1 (rPSGL-Ig) inhibits circulating activated platelet binding to neutrophils induced by damaged arterial surfaces. J Pharmacol Exp Ther. 298: 658-64 - Turner, C.P. <i>et al.</i> (2003) The role of P-selectin in the immune destruction of platelets. Br J

[Haematol. 121: 623-31.](#)

13. van Nispen tot Pannerden, H. *et al.* (2010) The platelet interior revisited: electron tomography reveals tubular alpha-granule subtypes. [Blood. 116: 1147-56.](#)

14. Knipe, L. *et al.* (2010) A revised model for the secretion of tPA and cytokines from cultured endothelial cells. [Blood. 116 \(12\): 2183-91.](#)

15. Kitaya, K. & Yasuo, T. (2010) Aberrant expression of selectin E, CXCL1, and CXCL13 in chronic endometritis. [Mod Pathol. 23 \(8\): 1136-46.](#)

16. Xiong, G.M. *et al.* (2015) Imparting electroactivity to polycaprolactone fibers with heparin-doped polypyrrole: Modulation of hemocompatibility and inflammatory responses. [Acta Biomater. 23: 240-9.](#)

17. Liao, Y. *et al.* (2017) Tailoring of TiO₂ films by H₂SO₄ treatment and UV irradiation to improve anticoagulant ability and endothelial cell compatibility. [Colloids Surf B Biointerfaces. 155: 314-22.](#)

Storage

Store at -20°C only.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life

18 months from date of despatch.

Health And Safety Information

Material Safety Datasheet documentation #10162 available at: 10162: <https://www.bio-rad-antibodies.com/uploads/MSDS/10162.pdf>

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87...)	Alk. Phos. , HRP
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®549 , DyLight®649 , DyLight®680 , DyLight®800 , FITC , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (STAR77...)	HRP
Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Rabbit Anti Mouse IgG (STAR8...)	DyLight®800
Goat Anti Mouse IgG (STAR70...)	FITC
Rabbit Anti Mouse IgG (STAR13...)	HRP
Human Anti Mouse IgG1 (HCA036...)	HRP

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South America

Tel: +1 800 265 7376
Fax: +1 919 878 3751
Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700
Fax: +44 (0)1865 852 739
Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21
Fax: +49 (0) 89 8090 95 50
Email: antibody_sales_de@bio-rad.com

Printed on 17 May 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)