

Datasheet: MCA812GA

Description:	MOUSE ANTI RABBIT IgM (B CELL MARKER)
Specificity:	IgM (B-CELL MARKER)
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	NRBM
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Rabbit
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0mg/ml
External Database Links	UniProt: P04221 Related reagents P03988 Related reagents

Fusion Partners	Spleen cells from immunised mice were fused with cells of the Mouse P3X63Ag8.653 myeloma cell line.
Specificity	<p>Mouse anti Rabbit IgM (B Cell Marker) antibody, clone NRBM specifically recognizes the μ chain of rabbit Ig.</p> <p>Mammalian IgM is produced and secreted by plasma cells located in bone marrow, lymph nodes and spleen. IgM is present in both a secreted polymeric form and as cell surface monomeric form on B cells.</p> <p>Clone NRBM has been used in number of studies to label IgM⁺ B cells (Dewals et al. 2011, Waclavicek et al. 2009) and as such can be considered a reliable marker of lagomorph (rabbit) B cells in flow cytometry.</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul.
References	<ol style="list-style-type: none"> 1. Idogawa, H. <i>et al.</i> (1997) Progression of articular destruction and the production of tumour necrosis factor-alpha in antigen-induced arthritis in rabbits. Scand J Immunol. 46 (6): 572-80. 2. Dewals, B. <i>et al.</i> (2008) Malignant catarrhal fever induced by alcelaphine herpesvirus 1 is associated with proliferation of CD8⁺ T cells supporting a latent infection. PLoS One 3: e1627. 3. Gillet, L. <i>et al.</i> (2009) Anchoring tick salivary anti-complement proteins IRAC I and IRAC II to membrane increases their immunogenicity. Vet Res. 40: 51. 4. Stich, N. <i>et al.</i> (2010) Staphylococcal Superantigen (TSST-1) Mutant Analysis Reveals that T Cell Activation Is Required for Biological Effects in the Rabbit Including the Cytokine Storm. Toxins 2: 2272-88. 5. Waclavicek, M. <i>et al.</i> (2009) Analysis of the early response to TSST-1 reveals Vbeta-unrestricted extravasation, compartmentalization of the response, and unresponsiveness but not anergy to TSST-1. J Leukoc Biol. 85 (1): 44-54. 6. Anderson, I.E. <i>et al.</i> (2008) Production and utilization of interleukin-15 in malignant catarrhal fever. J Comp Pathol. 138: 131-44. 7. Dewals, B.G. and Vanderplasschen, A. (2011) Malignant catarrhal fever induced by Alcelaphine herpesvirus 1 is characterized by an expansion of activated CD3⁺CD8⁺CD4⁻ T cells expressing a cytotoxic phenotype in both lymphoid and non-lymphoid tissues. Vet Res. 42: 95. 8. Dewals, B. <i>et al.</i> (2011) Ex vivo bioluminescence detection of alcelaphine herpesvirus 1 infection during malignant catarrhal fever. J Virol. 85: 6941-54. 9. Milanovic, V. <i>et al.</i> (2017) Histological and immunological changes in uterus during the different reproductive stages at Californian rabbit (<i>Oryctolagus cuniculus</i>). Kafkas Univ Vet Fak Derg. 23, 137-44.
Storage	<p>Store at +4°C or at -20°C if preferred.</p> <p>Storage in frost-free freezers is not recommended.</p> <p>This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.</p>
Shelf Life	18 months from date of despatch.
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: 10040: https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87...)	Alk. Phos. , HRP
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight@488 , DyLight@549 , DyLight@649 , DyLight@680 , DyLight@800 , FITC , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (STAR77...)	HRP
Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Rabbit Anti Mouse IgG (STAR8...)	DyLight@800
Goat Anti Mouse IgG (STAR70...)	FITC
Rabbit Anti Mouse IgG (STAR13...)	HRP
Human Anti Mouse IgG1 (HCA036...)	HRP

North & South America Tel: +1 800 265 7376
Fax: +1 919 878 3751
Email: antibody_sales_us@bio-rad.com

Worldwide Tel: +44 (0)1865 852 700
Fax: +44 (0)1865 852 739
Email: antibody_sales_uk@bio-rad.com

Europe Tel: +49 (0) 89 8090 95 21
Fax: +49 (0) 89 8090 95 50
Email: antibody_sales_de@bio-rad.com

'M316328:180503'

Printed on 05 May 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)