

Datasheet: MCA837GA

Description:	MOUSE ANTI BOVINE CD8
Specificity:	CD8
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	CC63
Isotype:	IgG2a
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			1/50
Immunohistology - Frozen	■			
Immunohistology - Paraffin	■			
ELISA			■	
Immunoprecipitation	■			
Western Blotting			■	
Immunofluorescence	■			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Bovine
Species Cross Reactivity	Reacts with: Sheep, Goat N.B. Antibody reactivity and working conditions may vary between species.
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1 mg/ml

**External Database
Links**

UniProt:

[P31783](#) [Related reagents](#)

Entrez Gene:

[281060](#) CD8A [Related reagents](#)

Fusion Partners Spleen cells from an immunised mouse were fused with cells of the mouse NS1 myeloma cell line.

Specificity **Mouse anti Bovine CD8 antibody, clone CC63** reacts with the bovine CD8 antigen expressed by a subset of T lymphocytes. The antibody precipitates molecules of ~34 kDa and ~38 kDa under reducing conditions. Clone CC63 has been reported as being suitable for use on formalin dichromate (FD5) fixed paraffin embedded tissue with amplification and antigen retrieval techniques ([Gutierrez *et al.* 1999](#)).

Flow Cytometry Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.

References

1. MacHugh, N.D. & Sopp P (1991) Individual antigens of cattle. Bovine CD8 (BoCD8). [Vet Immunol Immunopathol. 27 \(1-3\): 65-9.](#)
2. Gutierrez, M. *et al.* (1999) The detection of CD2+, CD4+, CD8+, and WC1+ T lymphocytes, B cells and macrophages in fixed and paraffin embedded bovine tissue using a range of antigen recovery and signal amplification techniques. [Vet Immunol Immunopathol. 71 \(3-4\): 321-34.](#)
3. Twizere, J.C. *et al.* (2000) Discordance between bovine leukemia virus tax immortalization *in vitro* and oncogenicity *in vivo*. [J Virol. 74 \(21\): 9895-902.](#)
4. Winkler, M.T. *et al.* (1999) Bovine herpesvirus 1 can infect CD4(+) T lymphocytes and induce programmed cell death during acute infection of cattle. [J Virol. 73 \(10\): 8657-68.](#)
5. Winkler, M.T. *et al.* (2000) Persistence and reactivation of bovine herpesvirus 1 in the tonsils of latently infected calves. [J Virol. 74 \(11\): 5337-46.](#)
6. Sidders, B. *et al.* (2008) Screening of highly expressed mycobacterial genes identifies Rv3615c as a useful differential diagnostic antigen for the *Mycobacterium tuberculosis* complex. [Infect Immun. 76: 3932-9.](#)
7. Sanchez, J. *et al.* (2011) Microscopical and immunological features of tuberculoid granulomata and cavitary pulmonary tuberculosis in naturally infected goats. [J Comp Pathol. 145 \(2-3\): 107-17.](#)
8. La Manna, M.P. *et al.* (2011) Expansion of intracellular IFN-γ positive lymphocytes during *Mycoplasma agalactiae* infection in sheep. [Res Vet Sci. 91 \(3\): e64-7.](#)
9. Fulton, B.E. Jr. *et al.* (2006) Dissemination of bovine leukemia virus-infected cells from a newly infected sheep lymph node. [J Virol. 80: 7873-84.](#)
10. Harris, J. *et al.* (2002) Expression of caveolin by bovine lymphocytes and antigen-presenting cells. [Immunology. 105: 190-5.](#)
11. Liebana, E. *et al.* (2007) Distribution and activation of T-lymphocyte subsets in tuberculous bovine lymph-node granulomas. [Vet Pathol. 44: 366-72.](#)
12. Lynch, E.M. *et al.* (2010) Effect of abrupt weaning at housing on leukocyte distribution, functional activity of neutrophils, and acute phase protein response of beef calves. [BMC Vet Res. 6: 39.](#)
13. Lacroux, C. *et al.* (2012) Prionemia and leukocyte-platelet-associated infectivity in sheep transmissible spongiform encephalopathy models. [J Virol. 86 \(4\): 2056-66.](#)
14. Coad, M. *et al.* (2010) Repeat tuberculin skin testing leads to desensitisation in naturally infected tuberculous cattle which is associated with elevated interleukin-10 and decreased interleukin-1 beta responses. [Vet Res. 41: 14.](#)
15. Toman, M. *et al.* (2003) Immunological characteristics of cattle with *Mycobacterium avium* subsp. *paratuberculosis* infection [Vet. Med. – Czech, 48, 2003: 147–54](#)
16. Constantinoiu, C.C. *et al.* (2010) Local immune response against larvae of *Rhipicephalus (Boophilus) microplus* in *Bos taurus indicus* and *Bos taurus taurus* cattle. [Int J Parasitol. 40:](#)

[865-75.](#)

17. Brodzki, P. *et al.* (2014) Phenotyping of leukocytes and granulocyte and monocyte phagocytic activity in the peripheral blood and uterus of cows with endometritis. [Theriogenology. 82 \(3\): 403-10.](#)
18. Romero-Palomo F *et al.* (2015) Immunopathologic Changes in the Thymus of Calves Pre-infected with BVDV and Challenged with BHV-1. [Transbound Emerg Dis. Aug 25. \[Epub ahead of print\]](#)
19. Stenfeldt, C. *et al.* (2015) Pathogenesis of Primary Foot-and-Mouth Disease Virus Infection in the Nasopharynx of Vaccinated and Non-Vaccinated Cattle. [PLoS One. 10 \(11\): e0143666.](#)
20. Leite FL *et al.* (2015) ZAP-70, CTLA-4 and proximal T cell receptor signaling in cows infected with *Mycobacterium avium* subsp. *paratuberculosis*. [Vet Immunol Immunopathol. 167 \(1-2\): 15-21.](#)
21. Silva, A.P. *et al.* (2015) Encapsulated *Brucella ovis* Lacking a Putative ATP-Binding Cassette Transporter (ΔabcBA) Protects against Wild Type *Brucella ovis* in Rams. [PLoS One. 10 \(8\): e0136865.](#)
22. Schmidt, N. *et al.* (2018) Decreased STEC shedding by cattle following passive and active vaccination based on recombinant *Escherichia coli* Shiga toxoids. [Vet Res. 49 \(1\): 28.](#)

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life

18 months from date of despatch.

Health And Safety Information

Material Safety Datasheet documentation #10040 available at: 10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87...)	Alk. Phos. , HRP
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®549 , DyLight®649 , DyLight®680 , DyLight®800 , FITC , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (STAR77...)	HRP
Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Rabbit Anti Mouse IgG (STAR8...)	DyLight®800
Goat Anti Mouse IgG (STAR70...)	FITC
Human Anti Mouse IgG2a (HCA037...)	FITC , HRP
Rabbit Anti Mouse IgG (STAR13...)	HRP

Recommended Negative Controls

[MOUSE IgG2a NEGATIVE CONTROL \(MCA929\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com 'M316335:180503'	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

Printed on 05 May 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)