

Datasheet: MCA87GA

Description:	MOUSE ANTI HUMAN CD45
Specificity:	CD45
Other names:	LCA
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	F10-89-4
Isotype:	IgG2a
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			1/10 - 1/50
Immunohistology - Frozen	■			1/500 - 1/1000
Immunohistology - Paraffin (1)	■			1/100 - 1/200
ELISA			■	
Immunoprecipitation	■			
Western Blotting			■	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1) This product requires antigen retrieval using heat treatment prior to staining of paraffin sections.

EDTA pH8.0 is recommended for this purpose.

Target Species	Human
Species Cross Reactivity	Reacts with: Horse N.B. Antibody reactivity and working conditions may vary between species.
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide

Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Human T lymphocytes.
External Database Links	UniProt: P08575 Related reagents Entrez Gene: 5788 PTPRC Related reagents
Synonyms	CD45
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the mouse NS-1 myeloma cell line.
Specificity	Mouse anti Human CD45 antibody, clone F10-89-4 recognizes the human CD45 cell surface antigen, also known as the leucocyte common antigen (LCA). CD45 is a complex molecule existing in a number of isoforms. Antibodies recognising a common epitope on all of these isoforms are termed CD45 whilst those recognising only individual isoforms are termed CD45RA or CD45RO etc. Mouse anti Human CD45 antibody, clone F10-89-4 reacts with all forms of CD45 expressed by all haematopoietic cells, except erythrocytes, having a higher level of expression on lymphocytes than on granulocytes. Mouse anti Human CD45 antibody, clone F10-89-4 is routinely tested in flow cytometry on human peripheral blood leucocytes
Flow Cytometry	Use 10ul of the suggested working dilution to label 10^6 cells in 100ul.
Histology Positive Control Tissue	Human tonsil
References	1. Dalchau, R. <i>et al.</i> (1980) Monoclonal antibody to a human leukocyte-specific membrane glycoprotein probably homologous to the leukocyte-common (L-C) antigen of the rat. Eur J Immunol. 10 (10): 737-44. 2. Quenby, S <i>et al.</i> (1999) Pre-implantation endometrial leukocytes in women with recurrent miscarriage. Human Reprod. 14(9):2386-2391. 3. Hauser, P.V. <i>et al.</i> (2010) Stem cells derived from human amniotic fluid contribute to acute kidney injury recovery. Am J Pathol. 177: 2011-21. 4. Mallam, E. <i>et al.</i> (2010) Characterization of <i>in vitro</i> expanded bone marrow-derived mesenchymal stem cells from patients with multiple sclerosis. Mult Scler. 16: 909-18. 5. Marrinucci, D. <i>et al.</i> (2010) Cytomorphology of circulating colorectal tumor cells:a small case series. J Oncol. 2010: 861341. 6. Kazane, S.A. <i>et al.</i> (2012) Site-specific DNA-antibody conjugates for specific and sensitive immuno-PCR. Proc Natl Acad Sci U S A. 109: 3731-6. 7. Paul, G. <i>et al.</i> (2012) The adult human brain harbors multipotent perivascular mesenchymal stem cells. PLoS One. 7: e35577.

8. Sadarangani, A. *et al.* (2015) GLI2 inhibition abrogates human leukemia stem cell dormancy. [J Transl Med. 13: 98.](#)
9. Gunawardene, P. *et al.* (2015) Association Between Circulating Osteogenic Progenitor Cells and Disability and Frailty in Older Persons: The Nepean Osteoporosis and Frailty Study. [J Gerontol A Biol Sci Med Sci. pii: glv190.](#)
10. Gogoi P *et al.* (2016) Development of an Automated and Sensitive Microfluidic Device for Capturing and Characterizing Circulating Tumor Cells (CTCs) from Clinical Blood Samples. [PLoS One. 11 \(1\): e0147400.](#)
11. Spaas, J.H. *et al.* (2013) Culture and characterisation of equine peripheral blood mesenchymal stromal cells. [Vet J. 195 \(1\): 107-13.](#)
12. Gomiero, C. *et al.* (2016) Tenogenic induction of equine mesenchymal stem cells by means of growth factors and low-level laser technology. [Vet Res Commun. 40 \(1\): 39-48.](#)
13. De Schauwer, C. *et al.* (2012) In search for cross-reactivity to immunophenotype equine mesenchymal stromal cells by multicolor flow cytometry. [Cytometry A. 81 \(4\): 312-23.](#)
14. Bianchessi, M. *et al.* (2016) Effect of Fibroblast Growth Factor 2 on Equine Synovial Fluid Chondroprogenitor Expansion and Chondrogenesis. [Stem Cells Int. 2016: 9364974.](#)
15. Mohamed Suhaimi, N.A. *et al.* (2015) Non-invasive sensitive detection of KRAS and BRAF mutation in circulating tumor cells of colorectal cancer patients. [Mol Oncol. 9 \(4\): 850-60.](#)
16. Ruiz, C. *et al.* (2015) Limited genomic heterogeneity of circulating melanoma cells in advanced stage patients. [Phys Biol. 12 \(1\): 016008.](#)
17. Branly, T. *et al.* (2017) Characterization and use of Equine Bone Marrow Mesenchymal Stem Cells in Equine Cartilage Engineering. Study of their Hyaline Cartilage Forming Potential when Cultured under Hypoxia within a Biomaterial in the Presence of BMP-2 and TGF- β 1. [Stem Cell Rev. Jun 09 \[Epub ahead of print\].](#)
18. GarikipatiV, N.S. *et al.* (2018) Isolation and characterization of mesenchymal stem cells from human fetus heart. [PLoS One. 13 \(2\): e0192244.](#)

Storage	Store at +4°C or at -20°C if preferred. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Shelf Life	18 months from date of despatch.
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: 10040: https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87...)	Alk. Phos. , HRP
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®549 , DyLight®649 , DyLight®680 , DyLight®800 , FITC , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (STAR77...)	HRP
Rabbit Anti Mouse IgG (STAR12...)	RPE

Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)
Rabbit Anti Mouse IgG (STAR8...) [DyLight®800](#)
Goat Anti Mouse IgG (STAR70...) [FITC](#)
Human Anti Mouse IgG2a (HCA037...) [FITC](#), [HRP](#)
Rabbit Anti Mouse IgG (STAR13...) [HRP](#)

Recommended Negative Controls

[MOUSE IgG2a NEGATIVE CONTROL \(MCA929\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

'M316343:180503'

Printed on 17 May 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)