

Datasheet: MCA87PBT

Description:	MOUSE ANTI HUMAN CD45:Pacific Blue®
Specificity:	CD45
Other names:	LCA
Format:	Pacific Blue®
Product Type:	Monoclonal Antibody
Clone:	F10-89-4
Isotype:	IgG2a
Quantity:	25 TESTS/0.25ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human		
Species Cross Reactivity	Reacts with: Horse N.B. Antibody reactivity and working conditions may vary between species.		
Product Form	Purified IgG conjugated to Pacific Blue® - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	Pacific Blue®	410	455
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative Stabilisers	0.09% Sodium Azide		
	1% Bovine Serum Albumin		
Approx. Protein Concentrations	IgG concentration 0.05 mg/ml		
Immunogen	Human T lymphocytes.		

**External Database
Links**

UniProt:

[P08575](#) [Related reagents](#)

Entrez Gene:

[5788](#) PTPRC [Related reagents](#)

Synonyms

CD45

Fusion Partners

Spleen cells from immunised BALB/c mice were fused with cells of the mouse NS-1 myeloma cell line.

Specificity

Mouse anti Human CD45 antibody, clone F10-89-4 recognizes the human CD45 cell surface antigen, also known as the leucocyte common antigen (LCA). CD45 is a complex molecule existing in a number of isoforms.

Antibodies recognising a common epitope on all of these isoforms are termed CD45 whilst those recognising only individual isoforms are termed CD45RA or CD45RO etc.

Mouse anti Human CD45 antibody, clone F10-89-4 reacts with all forms of CD45 expressed by all haematopoietic cells, except erythrocytes, having a higher level of expression on lymphocytes than on granulocytes.

Mouse anti Human CD45 antibody, clone F10-89-4 is routinely tested in flow cytometry on human peripheral blood leucocytes

Flow Cytometry

Use 10ul of the suggested working dilution to label 10^6 cells or 100ul whole blood.

References

1. Dalchau, R. *et al.* (1980) Monoclonal antibody to a human leukocyte-specific membrane glycoprotein probably homologous to the leukocyte-common (L-C) antigen of the rat. [Eur J Immunol. 10 \(10\): 737-44.](#)
2. Quenby, S *et al.* (1999) Pre-implantation endometrial leukocytes in women with recurrent miscarriage. [Human Reprod. 14\(9\):2386-2391.](#)
3. Hauser, P.V. *et al.* (2010) Stem cells derived from human amniotic fluid contribute to acute kidney injury recovery. [Am J Pathol. 177: 2011-21.](#)
4. Mallam, E. *et al.* (2010) Characterization of *in vitro* expanded bone marrow-derived mesenchymal stem cells from patients with multiple sclerosis. [Mult Scler. 16: 909-18.](#)
5. Marrinucci, D. *et al.* (2010) Cytomorphology of circulating colorectal tumor cells:a small case series. [J Oncol. 2010: 861341.](#)
6. Kazane, S.A. *et al.* (2012) Site-specific DNA-antibody conjugates for specific and sensitive immuno-PCR. [Proc Natl Acad Sci U S A. 109: 3731-6.](#)
7. Paul, G. *et al.* (2012) The adult human brain harbors multipotent perivascular mesenchymal stem cells. [PLoS One. 7: e35577.](#)
8. Sadarangani, A. *et al.* (2015) GLI2 inhibition abrogates human leukemia stem cell dormancy. [J Transl Med. 13: 98.](#)
9. Gunawardene, P. *et al.* (2015) Association Between Circulating Osteogenic Progenitor Cells and Disability and Frailty in Older Persons: The Nepean Osteoporosis and Frailty Study. [J Gerontol A Biol Sci Med Sci. pii: glv190.](#)
10. Gogoi P *et al.* (2016) Development of an Automated and Sensitive Microfluidic Device for Capturing and Characterizing Circulating Tumor Cells (CTCs) from Clinical Blood Samples. [PLoS One. 11 \(1\): e0147400.](#)
11. Spaas, J.H. *et al.* (2013) Culture and characterisation of equine peripheral blood mesenchymal stromal cells. [Vet J. 195 \(1\): 107-13.](#)

12. Gomiero, C. *et al.* (2016) Tenogenic induction of equine mesenchymal stem cells by means of growth factors and low-level laser technology. [Vet Res Commun. 40 \(1\): 39-48.](#)
13. De Schauwer, C. *et al.* (2012) In search for cross-reactivity to immunophenotype equine mesenchymal stromal cells by multicolor flow cytometry. [Cytometry A. 81 \(4\): 312-23.](#)
14. Bianchessi, M. *et al.* (2016) Effect of Fibroblast Growth Factor 2 on Equine Synovial Fluid Chondroprogenitor Expansion and Chondrogenesis. [Stem Cells Int. 2016: 9364974.](#)
15. Mohamed Suhaimi, N.A. *et al.* (2015) Non-invasive sensitive detection of KRAS and BRAF mutation in circulating tumor cells of colorectal cancer patients. [Mol Oncol. 9 \(4\): 850-60.](#)
16. Ruiz, C. *et al.* (2015) Limited genomic heterogeneity of circulating melanoma cells in advanced stage patients. [Phys Biol. 12 \(1\): 016008.](#)
17. Branly, T. *et al.* (2017) Characterization and use of Equine Bone Marrow Mesenchymal Stem Cells in Equine Cartilage Engineering. Study of their Hyaline Cartilage Forming Potential when Cultured under Hypoxia within a Biomaterial in the Presence of BMP-2 and TGF- β 1. [Stem Cell Rev. Jun 09 \[Epub ahead of print\].](#)
18. GarikipatiV, N.S. *et al.* (2018) Isolation and characterization of mesenchymal stem cells from human fetus heart. [PLoS One. 13 \(2\): e0192244.](#)

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life

18 months from date of despatch.

Acknowledgements

This product is provided under an intellectual property licence from Life Technologies Corporation. The transfer of this product is contingent on the buyer using the purchased product solely in research, excluding contract research or any fee for service research, and the buyer must not sell or otherwise transfer this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; (c) manufacturing or quality assurance or quality control, or (d) resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad CA 92008 USA or outlicensing@thermofisher.com

Health And Safety Information

Material Safety Datasheet documentation #10041 available at: 10041: <https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf>

Regulatory

For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG2a NEGATIVE CONTROL:Pacific Blue® \(MCA929PB\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M291309:160715'

Printed on 17 May 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)