

Datasheet: MCA90F

Description:	MOUSE ANTI HUMAN CD90:FITC
Specificity:	CD90
Other names:	THY1
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	F15-42-1
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. The suggested working dilution is given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human		
Species Cross Reactivity	Reacts with: Cynomolgus monkey N.B. Antibody reactivity and working conditions may vary between species.		
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525
Preparation	Purified IgG prepared by affinity chromatography on Protein G		
Buffer Solution	Phosphate buffered saline		
Preservative Stabilisers	0.09% Sodium Azide		
	1% Bovine Serum Albumin		
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml		
Immunogen	Purified human brain Thy-1.		

**External Database
Links**

UniProt:

[P04216](#) [Related reagents](#)

Entrez Gene:

[7070](#) THY1 [Related reagents](#)

Fusion Partners Spleen cells from immunised BALB/c mice were fused with cells of the mouse NS-1 myeloma cell line.

Specificity **Mouse anti Human CD90 antibody, clone F15-42-1** recognizes the human CD90 cell surface antigen, a ~25 kDa glycoprotein homologous to rat Thy1. The antigen is expressed by a subset of CD34+ve cells in the bone marrow and by prothymocytes within the thymus. CD90 is also expressed extensively within the brain.

Mouse anti Human CD90 antibody, clone F15-42-1 is routinely tested in flow cytometry on the MOLT4 cell line.

Flow Cytometry Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.

- References**
1. McKenzie, J.L. & Fabre, J.W. (1981) Human thy-1: unusual localization and possible functional significance in lymphoid tissues. [J Immunol. 126 \(3\): 843-50.](#)
 2. Daar, A.S. & Fabre, J.W. (1981) Demonstration with monoclonal antibodies of an unusual mononuclear cell infiltrate and loss of normal epithelial membrane antigens in human breast carcinomas. [Lancet. 2 \(8244\): 434-8.](#)
 3. Paul, G. *et al.* (2012) The adult human brain harbors multipotent perivascular mesenchymal stem cells. [PLoS One. 7: e35577.](#)
 4. Fiegel, H.C. *et al.* (2004) Stem-like cells in human hepatoblastoma. [J Histochem Cytochem. 52 \(11\): 1495-501.](#)
 5. Hagood, J.S. *et al.* (2005) Loss of fibroblast Thy-1 expression correlates with lung fibrogenesis. [Am J Pathol. 167 \(2\): 365-79.](#)
 6. Diaz-Romero, J. *et al.* (2008) Immunophenotypic changes of human articular chondrocytes during monolayer culture reflect bona fide dedifferentiation rather than amplification of progenitor cells. [J Cell Physiol. 214: 75-83.](#)
 7. Cox, G. *et al.* (2011) The use of the reamer-irrigator-aspirator to harvest mesenchymal stem cells. [J Bone Joint Surg Br. 93: 517-24.](#)
 8. Cizeau, J. *et al.* (2011) Fusogenics: a recombinant immunotoxin-based screening platform to select internalizing tumor-specific antibody fragments. [J Biomol Screen. 16: 90-100.](#)
 9. Gieseke, F. *et al.* (2010) Human multipotent mesenchymal stromal cells use galectin-1 to inhibit immune effector cells. [Blood. 116: 3770-9.](#)
 10. Hauser, P.V. *et al.* (2010) Stem cells derived from human amniotic fluid contribute to acute kidney injury recovery. [Am J Pathol. 177: 2011-21.](#)
 11. Holzwarth, C. *et al.* (2010) Low physiologic oxygen tensions reduce proliferation and differentiation of human multipotent mesenchymal stromal cells. [BMC Cell Biol. 11:11](#)
 12. Karlsen, T.A. *et al.* (2010) Human primary articular chondrocytes, chondroblasts-like cells, and dedifferentiated chondrocytes: differences in gene, microRNA, and protein expression and phenotype. [Tissue Eng Part C Methods. 17: 219-27.](#)
 13. Manochantr, S. *et al.* (2010) Isolation, characterization and neural differentiation potential of amnion derived mesenchymal stem cells. [J Med Assoc Thai. 93 Suppl 7: S183-91.](#)
 14. Meng, J. *et al.* (2011) Contribution of human muscle-derived cells to skeletal muscle regeneration in dystrophic host mice. [PLoS One. 6: e17454.](#)
 15. Pessina, A. *et al.* (2010) CD45+/CD133+ positive cells expanded from umbilical cord blood expressing PDX-1 and markers of pluripotency. [Cell Biol Int. 34: 783-90.](#)

16. Tome, M. *et al.* (2007) Calponin is expressed by subpopulations of connective tissue cells but not olfactory ensheathing cells in the neonatal olfactory mucosa. [BMC Neurosci. 8: 74.](#)
17. Yin, S. *et al.* (2010) Chondrogenic transdifferentiation of human dermal fibroblasts stimulated with cartilage-derived morphogenetic protein 1. [Tissue Eng Part A. 16: 1633-43.](#)
18. Shafaei, H. *et al.* (2011) Effects of human placental serum on proliferation and morphology of human adipose tissue-derived stem cells. [Bone Marrow Transplant. 46: 1464-71.](#)
19. Escobar, C.H. & Chaparro, O. (2016) Xeno-Free Extraction, Culture, and Cryopreservation of Human Adipose-Derived Mesenchymal Stem Cells. [Stem Cells Transl Med. 5 \(3\): 358-65.](#)
20. Shinoda, K. *et al.* (2016) Thy1+IL-7+ lymphatic endothelial cells in iBALT provide a survival niche for memory T-helper cells in allergic airway inflammation. [Proc Natl Acad Sci U S A. May 2. pii: 201512600. \[Epub ahead of print\]](#)
21. Kamprom, W. *et al.* (2016) Endothelial Progenitor Cell Migration-Enhancing Factors in the Secretome of Placental-Derived Mesenchymal Stem Cells. [Stem Cells Int. 2016: 2514326.](#)
22. Vaquero, J. *et al.* (2016) An approach to personalized cell therapy in chronic complete paraplegia: The Puerta de Hierro phase I/II clinical trial. [Cytotherapy. 18 \(8\): 1025-36.](#)
23. Zhang, X. *et al.* (2017) Regeneration of hyaline-like cartilage in situ with SOX9 stimulation of bone marrow-derived mesenchymal stem cells. [PLoS One. 12 \(6\): e0180138.](#)
24. GarikipatiV, N.S. *et al.* (2018) Isolation and characterization of mesenchymal stem cells from human fetus heart. [PLoS One. 13 \(2\): e0192244.](#)

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life

18 months from date of despatch.

Health And Safety Information

Material Safety Datasheet documentation #10041 available at:
10041: <https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf>

Regulatory

For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA928F\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South America

Tel: +1 800 265 7376

Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M301828:170109'

