

Datasheet: MCA928EL

Description:	MOUSE IgG1 NEGATIVE CONTROL:Low Endotoxin
Specificity:	MOUSE IgG1 NEGATIVE CONTROL
Format:	Low Endotoxin
Product Type:	Negative/Isotype Control
Isotype:	IgG1
Quantity:	0.5 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			*
Immunohistology - Frozen	■			
Immunohistology - Paraffin			■	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. * It is recommended that the user titrates the antibody for use in their own system to a concentration equivalent to their test reagents.

Target Species	Negative Control
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	None present
Carrier Free	Yes
Endotoxin Level	<0.01 EU/ug
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml

Specificity

Mouse IgG1 negative control is negative by flow cytometry on all human cells and cell lines tested. Further tests have also shown that this reagent is also suitable for use as a negative control with bovine ([Maslanka et al, 2012](#)), ovine, porcine ([Kapetanovic et al, 2012](#)), equine ([Jacks et al, 2007](#)), canine ([Maiolini et al, 2012](#)), lapine ([Pakandl et al, 2008](#)) and guinea-pig tissues.

This reagent recognises a rat cell surface marker, and therefore cannot be used as a negative control in this species.

Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	1. Kupatt, C. <i>et al.</i> (2000) c7E3Fab reduces postischemic leukocyte-thrombocyte interaction mediated by fibrinogen. Implications for myocardial reperfusion injury. Arterioscler Thromb Vasc Biol. 20 (10): 2226-32. 2. Dalli, J. <i>et al.</i> (2008) Annexin 1 mediates the rapid anti-inflammatory effects of neutrophil-derived microparticles. Blood. 112 (6): 2512-9. 3. Barratt-Due, A. <i>et al.</i> (2011) Ornithodoros moubata Complement Inhibitor Is an Equally Effective C5 Inhibitor in Pigs and Humans. J Immunol. 187: 4913-9. 4. Kapetanovic, R. <i>et al.</i> (2012) Pig bone marrow-derived macrophages resemble human macrophages in their response to bacterial lipopolysaccharide. J Immunol. 188: 3382-94. 5. Maiolini, A. <i>et al.</i> (2012) Toll-like receptors 4 and 9 are responsible for the maintenance of the inflammatory reaction in canine steroid-responsive meningitis-arteritis, a large animal model for neutrophilic meningitis. J Neuroinflammation. 9: 226. 6. Maślanka, T. <i>et al.</i> (2012) The presence of CD25 on bovine WC1+ γδ T cells is positively correlated with their production of IL-10 and TGF-β, but not IFN-γ Polish Journal of Veterinary Sciences.15: 11–20. 7. Pakandl, M. <i>et al.</i> (2008) Immune response to rabbit coccidiosis: a comparison between infections with Eimeria flavescens and E. intestinalis. Folia Parasitol (Praha). 55:1-6. 8. Jacks, S. <i>et al.</i> (2007) Experimental infection of neonatal foals with Rhodococcus equi triggers adult-like gamma interferon induction. Clin Vaccine Immunol.14:669-77 9. Kamble, N.M. <i>et al.</i> (2016) Interaction of a live attenuated <i>Salmonella gallinarum</i> vaccine candidate with chicken bone marrow-derived dendritic cells. Avian Pathol. Jan 26:1-24. [Epub ahead of print] 10. Brace, P.T. <i>et al.</i> (2017) <i>Mycobacterium tuberculosis</i> subverts negative regulatory pathways in human macrophages to drive immunopathology. PLoS Pathog. 13 (6): e1006367. 11. Topoluk, N. <i>et al.</i> (2017) Amniotic Mesenchymal Stromal Cells Exhibit Preferential Osteogenic and Chondrogenic Differentiation and Enhanced Matrix Production Compared With Adipose Mesenchymal Stromal Cells. Am J Sports Med. 363546517706138. 12. Iwaszko-Simonik, A. <i>et al.</i> (2015) Expression of surface platelet receptors (CD62P and CD41/61) in horses with recurrent airway obstruction (RAO). Vet Immunol Immunopathol. 164 (1-2): 87-92. 13. Arzi, B. <i>et al.</i> (2017) Therapeutic Efficacy of Fresh, Allogeneic Mesenchymal Stem Cells for Severe Refractory Feline Chronic Gingivostomatitis. Stem Cells Transl Med. 6 (8): 1710-22.
Storage	Store at -20°C only. This product should be stored undiluted. Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Shelf Life	18 months from date of despatch.
Health And Safety Information	Material Safety Datasheet documentation #10162 available at: 10162: https://www.bio-rad-antibodies.com/uploads/MSDS/10162.pdf
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL:Low Endotoxin (MCA1209EL)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

'M316373:180503'

Printed on 21 Jun 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)