

Datasheet: MCA929B

Description:	MOUSE IgG2a NEGATIVE CONTROL:Biotin
Specificity:	MOUSE IgG2a NEGATIVE CONTROL
Format:	Biotin
Product Type:	Negative/Isotype Control
Isotype:	IgG2a
Quantity:	100 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			*
Immunohistology - Frozen			■	
Immunohistology - Paraffin			■	
ELISA			■	
Western Blotting			■	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures.

*It is recommended that the user dilutes the antibody for use in their own system to a concentration equivalent to their test reagents.

Target Species	Negative Control
Product Form	Purified IgG conjugated to Biotin - liquid
Preparation	Purified IgG prepared by affinity chromatography
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide 1% Bovine Serum Albumin
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml
Immunogen	Activated rat T-helper cells.
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the NS1 mouse myeloma cell line.
Specificity	Mouse IgG2a negative control antibody, clone OX-34 is suitable for use as a negative control reagent for the measurement of non-specific binding of mouse monoclonal antibodies of isotype

IgG2a to human tissue.

Clone MRC OX-34 recognises a rat cell surface marker, and therefore cannot be used as a negative control in this species.

This product is routinely tested in flow cytometry on rat splenocytes to confirm antibody activity and on human whole blood to test for suitability as a negative control.

Test results have shown that MCA929 is also suitable for use as a negative control with bovine, ovine, porcine, equine, canine, lapine and guinea-pig tissues.

This antibody may not be suitable for intracellular staining on some cell types.

Flow Cytometry	Use 10ul of the suggested working dilution to label 10^6 in 100ul.
References	1. Avigdor, A. <i>et al.</i> (2004) CD44 and hyaluronic acid cooperate with SDF-1 in the trafficking of human CD34+ stem/progenitor cells to bone marrow. Blood. 103 (8): 2981-9. 2. Kamble, N.M. <i>et al.</i> (2016) Interaction of a live attenuated <i>Salmonella gallinarum</i> vaccine candidate with chicken bone marrow-derived dendritic cells. Avian Pathol. Jan 26:1-24. [Epub ahead of print] 3. Wattegedera, S.R. <i>et al.</i> (2017) Enhancing the toolbox to study IL-17A in cattle and sheep. Vet Res. 48 (1): 20.
Storage	Store at +4°C or at -20°C if preferred. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Shelf Life	18 months from date of despatch.
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: 10041: https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf
Regulatory	For research purposes only

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M281126:160211'

Printed on 01 Aug 2018