

Datasheet: OBT1388

Description:	GOAT ANTI HUMAN ANNEXIN II (N-TERMINAL)
Specificity:	ANNEXIN II (N-TERMINAL)
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
ELISA	▪			1/4000
Western Blotting	▪			1 ug/ml - 3 ug/ml

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human
Species Cross Reactivity	Reacts with: Mouse Based on sequence similarity, is expected to react with: Rat, Bovine, Pig N.B. Antibody reactivity and working conditions may vary between species.
Product Form	Purified IgG - liquid
Antiserum Preparation	Antiserum to human Annexin II was raised by repeated immunisation of goats with highly purified antigen. Purified IgG prepared by affinity chromatography
Buffer Solution	TRIS buffered saline
Preservative	0.02% Sodium Azide (NaN ₃)
Stabilisers	0.5% Bovine Serum Albumin
Approx. Protein Concentrations	IgG concentration 0.5mg/ml
Immunogen	Peptide sequence STVHEILCKLSLEGD corresponding to the N-terminal region of Annexin II (NP_001002858.1; NP_004030.1; NP_001002857.1).
External Database Links	UniProt: P07355 Related reagents

Entrez Gene:[302](#) ANXA2 [Related reagents](#)

Synonyms	ANX2, ANX2L4, CAL1H, LPC2D
-----------------	----------------------------

Specificity	Goat anti Human Annexin II (N-Terminal) antibody recognizes an epitope within the N terminal (NT) region of Annexin II, a member of a family of structurally related calcium binding proteins implicated in mediating the the intracellular action of calcium. Cellular levels of annexin II mRNA and protein are strictly regulated during the mammalian cell cycle with maximal expression as cells enter S-phase.
--------------------	---

Western Blotting	OBT1388 detects a band of approximately 35 kDa in A431 and A549 cell lysates
-------------------------	--

References	1. Xie, R. <i>et al.</i> (2012) Phagocytosis by macrophages and endothelial cells inhibits procoagulant and fibrinolytic activity of acute promyelocytic leukemia cells. Blood. 119 (10): 2325-34.
-------------------	--

Storage	Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
----------------	---

Shelf Life	18 months from date of despatch.
-------------------	----------------------------------

Health And Safety Information	Material Safety Datasheet documentation #10058 available at: 10058: https://www.bio-rad-antibodies.com/uploads/MSDS/10058.pdf
--------------------------------------	--

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Secondary Antibodies

Rabbit Anti Goat IgG (Fc) (STAR122...) [FITC](#), [HRP](#)**North & South** Tel: +1 800 265 7376**America** Fax: +1 919 878 3751Email: antibody_sales_us@bio-rad.com**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M282562:160222'

Printed on 30 Apr 2018