

Datasheet: STAR104D488GA

Description:	GOAT F(ab') ₂ ANTI HAMSTER IgG:DyLight®488
Specificity:	IgG
Format:	DyLight®488
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			
Immunofluorescence	■			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls

Target Species	Hamster		
Product Form	F(ab') ₂ fragment of purified IgG conjugated to DyLight®488 - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	Dylight®488	493	518

Antiserum Preparation Antisera to hamster IgG were raised by repeated immunisation of goats with highly purified antigen. Purified IgG was prepared by affinity chromatography

Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Hamster IgG
Specificity	Goat F(ab')₂ anti Hamster IgG antibody recognizes hamster IgG. Goat F(ab') ₂ anti Hamster IgG antibody has been adsorbed against both mouse and rat immunoglobulins to minimise cross-reactivity.
Flow Cytometry	Use 50ul of the suggested working dilution to label 1 x 10 ⁶ cells in 100ul

References	1. Osorio, Y. <i>et al.</i> (2011) Identification of small molecule lead compounds for visceral leishmaniasis using a novel <i>ex vivo</i> splenic explant model system. PLoS Negl Trop Dis. 5 (2): e962. 2. Bouma, G. <i>et al.</i> (2011) Cytoskeletal remodeling mediated by WASp in dendritic cells is necessary for normal immune synapse formation and T-cell priming. Blood. 118 (9): 2492-501.
Storage	Store at +4°C or at -20°C if preferred. This product should be stored undiluted. Storage in frost free freezers is not recommended. This product is photosensitive and should be protected from light. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use
Shelf Life	18 months from date of despatch
Acknowledgements	DyLight® is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: 10040: https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf
Regulatory	For research purposes only

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

'M314248:180412'

Printed on 20 Jun 2018