

Datasheet: STAR104F

Description:	GOAT F(ab') ₂ ANTI HAMSTER IgG:FITC
Specificity:	IgG
Format:	FITC
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.4 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/100 - 1/200
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Hamster		
Product Form	F(ab') ₂ fragment of purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525

Antiserum Preparation Antisera to hamster IgG were raised by repeated immunisation of goats with highly purified antigen. Purified IgG was prepared by affinity chromatography.

Buffer Solution	Phosphate buffered saline
Preservative	0.09% Sodium Azide
Stabilisers	0.5% Bovine Serum Albumin

Approx. Protein Concentrations	IgG concentration 0.8 mg/ml
---------------------------------------	-----------------------------

Immunogen	Hamster IgG
------------------	-------------

Specificity	Goat F(ab')₂ anti Hamster IgG antibody recognizes hamster IgG. Goat F(ab') ₂ anti Hamster IgG antibody has been adsorbed against both mouse and rat immunoglobulins to minimise cross-reactivity.
--------------------	--

Flow Cytometry Use 50ul of the suggested working dilution to label 1 x 10⁶ cells in 100ul.

References

1. Osorio, Y. *et al.* (2011) Identification of small molecule lead compounds for visceral leishmaniasis using a novel *ex vivo* splenic explant model system. [PLoS Negl Trop Dis. 5 \(2\): e962.](#)
2. Bouma, G. *et al.* (2011) Cytoskeletal remodeling mediated by WASp in dendritic cells is necessary for normal immune synapse formation and T-cell priming. [Blood. 118 \(9\): 2492-501.](#)

Storage Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life 18 months from date of despatch.

Health And Safety Information Material Safety Datasheet documentation #10041 available at:
10041: <https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf>

Regulatory For research purposes only

North & South America Tel: +1 800 265 7376
Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700
Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21
Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M316887:180612'

Printed on 20 Jun 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)