

Datasheet: STAR110

Description:	GOAT ANTI MOUSE IgE
Specificity:	IgE
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.5 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			■	
Immunohistology - Frozen			■	
Immunohistology - Paraffin			■	
ELISA	■			1/100 - 1/1000
Immunoprecipitation			■	
Western Blotting			■	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography
Antiserum Preparation	Antisera to mouse IgE were raised by repeated immunisations of goats with highly purified antigen.
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Native mouse IgE.

External Database Links

UniProt:

[P06336](#)

[Related reagents](#)

Entrez Gene:

[380792](#) Gm900 [Related reagents](#)

Specificity **Goat anti mouse IgE polyclonal antibody** recognizes mouse IgE and shows negligible cross-reactivity with other immunoglobulins.

References

1. Xia, Z.W. *et al.* (2007) Heme oxygenase-1 attenuates ovalbumin-induced airway inflammation by up-regulation of foxp3 T-regulatory cells, interleukin-10, and membrane-bound transforming growth factor-1. [Am J Pathol. 171: 1904-14.](#)
2. Hartmann, W. *et al.* (2011) A novel and divergent role of granzyme a and B in resistance to helminth infection. [J Immunol. 186: 2472-81.](#)
3. Hadzhieva, M. *et al.* (2015) Mechanism and functional implications of the heme-induced binding promiscuity of IgE. [Biochemistry. 54 \(11\): 2061-72.](#)
4. Su, W. *et al.* (2012) Synthesized OVA323-339MAP octamers mitigate OVA-induced airway inflammation by regulating Foxp3 T regulatory cells. [BMC Immunol. 13: 34.](#)
5. Wei, J.F. *et al.* (2009) Induction of mast cell accumulation, histamine release and skin edema by N49 phospholipase A2. [BMC Immunol. 10: 21.](#)

Storage Store at +4°C. DO NOT FREEZE.
This product should be stored undiluted. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life 12 months from date of despatch.

Health And Safety Information Material Safety Datasheet documentation available at:
Material Safety Datasheet Documentation #10040 available at:
<https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Goat IgG (Fc) (STAR122...) [FITC](#), [HRP](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M314193:180412'

Printed on 13 Apr 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)