

Datasheet: VMA00221

Description:	MOUSE ANTI Cdc25C
Specificity:	Cdc25C
Format:	Purified
Product Type:	PrecisionAb™ Monoclonal
Isotype:	IgG1
Quantity:	100 µl

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	▪			1/1000

PrecisionAb antibodies have been extensively [validated for the western blot application](#). The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependant on sample type.

Target Species	Human
Species Cross Reactivity	Reacts with: Mouse, Rat N.B. Antibody reactivity and working conditions may vary between species.
Product Form	Purified IgG - liquid
Preparation	Mouse monoclonal antibody prepared by affinity chromatography on Protein G from ascites
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Immunogen	Purified recombinant fragment of human Cdc25C expressed in <i>E. coli</i>
External Database Links	<p>UniProt: P30307 Related reagents</p> <p>Entrez Gene: 995 CDC25C Related reagents</p>

Specificity

Mouse anti Human Cdc25C antibody recognizes the M-phase inducer phosphatase 3, also known as M-phase inducer phosphatase 3, dual specificity phosphatase Cdc25C, mitosis inducer

CDC25, phosphotyrosine phosphatase, protein phosphatase 1 and regulatory subunit 60.

The CDC25C gene is highly conserved during evolution and it plays a key role in the regulation of cell division. The encoded protein is a tyrosine phosphatase and belongs to the Cdc25 phosphatase family. It directs dephosphorylation of cyclin B-bound CDC2 and triggers entry into mitosis. It is also thought to suppress p53-induced growth arrest. Multiple alternatively spliced transcript variants of CDC25C have been described, however, the full-length nature of many of them is not known (provided by RefSeq, Jul 2008).

Mouse anti Human Cdc25C antibody detects a band of 75 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.

Western Blotting	Anti Cdc25C detects a band of approximately 75 kDa in HEK293 cell lysates
Instructions For Use	Please refer to the PrecisionAb western blotting protocol . For additional information on secondary antibody dilution and exposure time see product web page.
Storage	Store undiluted at -20°C, avoiding repeated freeze thaw cycles.
Shelf Life	As supplied, 12 months from date of despatch.
Acknowledgements	PrecisionAb™ is a trademark of Bio-Rad Laboratories.
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: Antibody (10040): https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (H/L) (STAR207...) [HRP](#)

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M306945:170707'

Printed on 01 May 2018