

## Datasheet: VMA00401KT

| Description:  | IGF2BP-2 ANTIBODY WITH CONTROL LYSATE |  |  |  |  |
|---------------|---------------------------------------|--|--|--|--|
| Specificity:  | IGF2BP-2                              |  |  |  |  |
| Format:       | Purified                              |  |  |  |  |
| Product Type: | PrecisionAb™ Monoclonal               |  |  |  |  |
| Clone:        | OTI3C8                                |  |  |  |  |
| Isotype:      | lgG1                                  |  |  |  |  |
| Quantity:     | 2 Westerns                            |  |  |  |  |

# **Product Details**

| W                  | estern Blotting<br>recisionAb an   | s, please v     | isit <u>www.b</u><br>Yes | io-rad-antib<br>No | odies.com/protocols. |   |  |  |  |  |  |  |  |
|--------------------|--|-----------------|--------------------------|--------------------|----------------------|---|--|--|--|--|--|--|--|
| Pi                 | ecisionAb an   |                 | Yes                      | No                 |                      | recommendations, please visit www.bio-rad-antibodies.com/protocols. |  |  |  |  |  |  |  |
| Pi                 | ecisionAb an   |                 |                          | No                 | Not Determined       | Suggested Dilution  |  |  |  |  |  |  |  |
|                    |  |                 | •                        |                    |                      | 1/1000  |  |  |  |  |  |  |  |
| us                 | <b>PrecisionAb antibodies have been extensively</b> <u>validated for the western blot application</u> . The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependent on sample type. |                 |                          |                    |                      |   |  |  |  |  |  |  |  |
| Target Species H   | uman   |                 |                          |                    |                      |   |  |  |  |  |  |  |  |
| Product Form Pr    | Purified IgG - liquid  |                 |                          |                    |                      |   |  |  |  |  |  |  |  |
| Preparation 20     | 20µl Mouse monoclonal antibody purified by affinity chromatography from ascites  |                 |                          |                    |                      |   |  |  |  |  |  |  |  |
| Buffer Solution Pl | Phosphate buffered saline  |                 |                          |                    |                      |   |  |  |  |  |  |  |  |
| Preservative 0.    | 0.09% Sodium Azide (NaN <sub>3</sub> )   |                 |                          |                    |                      |   |  |  |  |  |  |  |  |
| Stabilisers 19     | 1% Bovine Serum Albumin  |                 |                          |                    |                      |   |  |  |  |  |  |  |  |
| 50                 | % Glycerol   |                 |                          |                    |                      |   |  |  |  |  |  |  |  |
| Immunogen Fu       | Full length recombinant human IGF2BP-2 (NP_006539) produced in HEK293T cells   |                 |                          |                    |                      |   |  |  |  |  |  |  |  |
| External Database  | niProt:  |                 |                          |                    |                      |   |  |  |  |  |  |  |  |
| Links              | <u>Q9Y6M1</u>  | Related rea     | agents                   |                    |                      |   |  |  |  |  |  |  |  |
| E                  | ntrez Gene:  |                 |                          |                    |                      |   |  |  |  |  |  |  |  |
|                    | <u>10644</u> IGF2  | BP2 <u>Rela</u> | ited reage               | <u>nts</u>         |                      |   |  |  |  |  |  |  |  |
| Synonyms IN        | IP2, VICKZ2  |                 |                          |                    |                      |   |  |  |  |  |  |  |  |

| Specificity                      | <b>Mouse anti Human IGF2BP-2 antibody</b> recognizes insulin-like growth factor 2 mRNA-binding protein 2 (IGFBP-2), also known as IGF2 binding protein 2 and VICKZ family member 2.   |  |  |  |  |
|----------------------------------|---|--|--|--|--|
|                                  | IGF2BP-2 is a member of the IGF-II mRNA-binding protein (IMP) family. It functions by binding to the 5' UTR of the insulin-like growth factor 2 (IGF2) mRNA and regulating IGF2 translation.<br>Alternate transcriptional splice variants, encoding different isoforms, have been characterized (provided by RefSeq, Jul 2008). |  |  |  |  |
|                                  | Mouse anti IGF2BP-2 antibody detects bands of 62 and 66 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.   |  |  |  |  |
| Western Blotting                 | Anti IGF2BP-2 detects a bands of approximately 62 and 66 kDa in HepG2 cell lysates  |  |  |  |  |
| Instructions For Use             | Please refer to the <u>PrecisionAb western blotting protocol</u> . For additional information on secondary antibody dilution and exposure time see product web page.  |  |  |  |  |
| Lysate Composition               | 400µg HepG2 lysate lyophilized in RIPA buffer   |  |  |  |  |
| Lysate Reconstitution            | - If using DDT reconstitute the lyophilized lysate with 190 $\mu$ l DI H <sub>2</sub> O, add 200 $\mu$ l 2x Laemmli Sample Buffer and 10 $\mu$ l 2M DTT.  |  |  |  |  |
|                                  | - If using BME reconstitute the lyophilized lysate with 180µl DI H <sub>2</sub> O, add 200µl 2x Laemmli Sample Buffer and 20µl BME.   |  |  |  |  |
|                                  | Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol.  |  |  |  |  |
| Storage                          | Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles.   |  |  |  |  |
|                                  | Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage.  |  |  |  |  |
| Shelf Life                       | As supplied, 12 months from date of despatch  |  |  |  |  |
| Acknowledgements                 | PrecisionAb <sup>™</sup> is a trademark of Bio-Rad Laboratories   |  |  |  |  |
| Health And Safety<br>Information | Material Safety Datasheet documentation #10048 #10561 available at:<br>Antibody (10048): <u>https://www.bio-rad-antibodies.com/uploads/MSDS/10048.pdf</u><br>Lysate Material (10561): <u>https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf</u>  |  |  |  |  |
| Regulatory                       | For research purposes only  |  |  |  |  |

### Related Products

### **Recommended Secondary Antibodies**

Goat Anti Mouse IgG (H/L) (STAR207...) HRP

#### **Recommended Negative Controls**

MOUSE IgG1 NEGATIVE CONTROL (MCA928)

| North & South | Tel: +1 800 265 7376             | Worldwide |
|---------------|----------------------------------|-----------|
| America       | Fax: +1 919 878 3751             |           |
|               | Email: antibody_sales_us@bio-rad | .com      |

 Tel: +44 (0)1865 852 700
 Europe

 Fax: +44 (0)1865 852 739
 Email: antibody\_sales\_uk@bio-rad.com

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody\_sales\_de@bio-rad.com

'M286448:160718'

© 2018 Bio-Rad Laboratories Inc | Legal | Imprint