

## Datasheet: VMA00124K

<b>Description:</b>	RAF-1 ANTIBODY WITH CONTROL LYSATE
<b>Specificity:</b>	RAF-1
<b>Format:</b>	Purified
<b>Product Type:</b>	PrecisionAb™ Monoclonal
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	10 Westerns

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	▪			1/1000

**PrecisionAb antibodies have been extensively [validated for the western blot application](#).**

The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependant on sample type.

<b>Target Species</b>	Human
<b>Species Cross Reactivity</b>	Reacts with: Mouse, Rat <b>N.B.</b> Antibody reactivity and working conditions may vary between species.
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	100µl Mouse monoclonal antibody prepared by affinity chromatography on Protein G from ascites
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide (NaN <sub>3</sub> )
<b>Immunogen</b>	Purified recombinant fragment of human RAF-1 expressed in <i>E. coli</i>
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">P04049</a>   <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">5894</a>   RAF1   <a href="#">Related reagents</a></p>

<b>Synonyms</b>	RAF
<b>Specificity</b>	<p><b>Mouse anti Human RAF-1 antibody</b> recognizes RAF-1, also known as C-Raf proto-oncogene, serine/threonine kinase, Oncogene RAF1, RAF proto-oncogene serine/threonine-protein kinase, proto-oncogene c-RAF, raf proto-oncogene serine/threonine protein kinase or v-raf-1 murine leukemia viral oncogene homolog 1.</p> <p>RAF-1 is the cellular homolog of viral raf gene (v-raf). The encoded protein is a MAP kinase kinase kinase (MAP3K), which functions downstream of the Ras family of membrane associated GTPases to which it binds directly. Once activated, the cellular RAF-1 protein can phosphorylate to activate the dual specificity protein kinases MEK1 and MEK2, which in turn phosphorylate to activate the serine/threonine specific protein kinases, ERK1 and ERK2. Activated ERKs are pleiotropic effectors of cell physiology and play an important role in the control of gene expression involved in the cell division cycle, apoptosis, cell differentiation and cell migration. Mutations in RAF-1 are associated with Noonan syndrome 5 and LEOPARD syndrome 2 (provided by RefSeq, Jul 2008).</p> <p>Mouse anti Human RAF-1 antibody detects a band of 73 kDa. The antibody has been extensively validated for western blotting using whole cell lysates. This antibody is available with a positive control lysate, in a smaller trial size, and as a standalone antibody.</p>
<b>Western Blotting</b>	Anti RAF-1 detects a band of approximately 73 kDa in HEK293 cell lysates.
<b>Instructions For Use</b>	Please refer to the <a href="#">PrecisionAb western blotting protocol</a> . For additional information on secondary antibody dilution and exposure time see product web page.
<b>Lysate Composition</b>	400µg HEK293 lysate lyophilized in RIPA buffer.
<b>Lysate Reconstitution</b>	<p>- If using DDT reconstitute the lyophilized lysate with 190µl DI H<sub>2</sub>O, add 200µl 2x Laemmli Sample Buffer and 10µl 2M DTT.</p> <p>- If using BME reconstitute the lyophilized lysate with 180µl DI H<sub>2</sub>O, add 200µl 2x Laemmli Sample Buffer and 20µl BME.</p> <p>Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol.</p>
<b>Storage</b>	<p>Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles.</p> <p>Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage.</p>
<b>Shelf Life</b>	As supplied, 12 months from date of despatch.
<b>Acknowledgements</b>	PrecisionAb™ is a trademark of Bio-Rad Laboratories.
<b>Health And Safety Information</b>	<p>Material Safety Datasheet Documentation #10040 for antibody available at <a href="https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf</a></p> <p>Lysate Material Safety Datasheet Documentation #10561 available at <a href="https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf</a></p>
<b>Regulatory</b>	For research purposes only

## Related Products

### Recommended Secondary Antibodies

Goat Anti Mouse IgG (H/L) (STAR207...)[HRP](#)

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

'M306919:170707'

**Printed on 07 Jul 2017**

---

© 2017 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)