Datasheet: VMA00576

Description:	MOUSE ANTI TAB1
Specificity:	TAB1
Format:	Purified
Product Type:	PrecisionAb™ Monoclonal
Isotype:	IgG2a
Quantity:	100 µl

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	▪			1/1000

PrecisionAb antibodies have been extensively validated for the western blot application. The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependant on sample type.

Target Species

Human

Species Cross Reactivity

Reacts with: Mouse
N.B. Antibody reactivity and working conditions may vary between species.

Product Form

Purified IgG - liquid

Preparation

Mouse monoclonal antibody purified by affinity chromatography on Protein G from ascites

Buffer Solution

HEPES buffered saline

Preservative Stabilisers

0.09% Sodium Azide (NaN_3)
1% Bovine Serum Albumin
<50% Glycerol

Immunogen

E. coli-derived recombinant human TAB1 (amino acids 1 - 300)

External Database Links

UniProt:
Q15750 Related reagents

Entrez Gene:
10454 TAB1 Related reagents

Synonyms

MAP3K7IP1

Specificity	**Mouse anti Human TAB1 antibody** recognizes the TGF-beta-activated kinase 1 and MAP3K7-binding protein 1, also known as TAK1-binding protein 1, mitogen-activated protein kinase kinase kinase 7-interacting protein 1 or transforming growth factor beta-activated kinase-binding protein 1. The protein encoded by TAB1 gene was identified as a regulator of the MAP kinase kinase kinase MAP3K7/TAK1, which is known to mediate various intracellular signaling pathways, such as those induced by TGF beta, interleukin 1, and WNT-1. This protein interacts and thus activates TAK1 kinase. It has been shown that the C-terminal portion of this protein is sufficient for binding and activation of TAK1, while a portion of the N-terminus acts as a dominant-negative inhibitor of TGF beta, suggesting that this protein may function as a mediator between TGF beta receptors and TAK1. This protein can also interact with and activate the mitogen-activated protein kinase 14 (MAPK14/p38alpha). Alternatively spliced transcript variants encoding distinct isoforms have been reported (provided by RefSeq, Jul 2008). Mouse anti Human TAB1 antibody detects a band of 60 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.
Western Blotting	Anti TAB1 detects a band of approximately 60 kDa in HEK293 cell lysates
Instructions For Use	Please refer to the PrecisionAb western blotting protocol. For additional information on secondary antibody dilution and exposure time see product web page.
Storage	Store undiluted at -20^{o}C, avoiding repeated freeze thaw cycles.
Shelf Life	As supplied, 12 months from date of despatch.
Acknowledgements	PrecisionAb™ is a trademark of Bio-Rad Laboratories.
Health And Safety Information	Material Safety Datasheet documentation #20359 available at: Antibody (20359): https://www.bio-rad-antibodies.com/uploads/MSDS/20359.pdf
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (H/L) (STAR207...) HRP

Recommended Negative Controls

MOUSE IgG2a NEGATIVE CONTROL (MCA929)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	**Worldwide**	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	**Europe**	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com

'M302219:170201'

Printed on 01 May 2018