

Datasheet: VPA00430

Description:	RABBIT ANTI CORTACTIN
Specificity:	CORTACTIN
Format:	Purified
Product Type:	PrecisionAb™ Polyclonal
Isotype:	Polyclonal IgG
Quantity:	100 µl

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	■			1/1000

PrecisionAb antibodies have been extensively [validated for the western blot application](#). The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependant on sample type.

Target Species	Human
Species Cross Reactivity	Reacts with: Rat N.B. Antibody reactivity and working conditions may vary between species.
Product Form	Purified IgG - liquid
Preparation	Rabbit polyclonal antibody purified by affinity chromatography
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Immunogen	KLH conjugated synthetic peptide corresponding to amino acid 273-302 of human cortactin
External Database Links	UniProt: Q14247 Related reagents Entrez Gene: 2017 CTTN Related reagents
Synonyms	EMS1

Specificity **Rabbit anti Human cortactin antibody** recognizes the Src substrate cortactin, also known as amplexin, ems1 sequence (mammary tumor and squamous cell carcinoma-associated (p80/85 src substrate) or oncogene EMS1.

CTTN is overexpressed in breast cancer and squamous cell carcinomas of the head and neck ([Patel et al. 1996](#)). Cortactin is localized in the cytoplasm and in areas of the cell-substratum contacts. CTTN has two roles: (1) regulating the interactions between components of adherens-type junctions and (2) organizing the cytoskeleton and cell adhesion structures of epithelia and carcinoma cells. During apoptosis, the encoded protein is degraded in a caspase-dependent manner. The aberrant regulation of CTTN contributes to tumor cell invasion and metastasis ([Li et al. 2001](#)). Three splice variants that encode different isoforms have been identified for CTTN (provided by RefSeq, May 2010).

Rabbit anti Human cortactin antibody detects a band of 80 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.

Western Blotting	Anti cortactin detects a band of approximately 80 kDa in A431 cell lysates
Instructions For Use	Please refer to the PrecisionAb western blotting protocol . For additional information on secondary antibody dilution and exposure time see product web page.
Storage	Store undiluted at -20°C, avoiding repeated freeze thaw cycles
Shelf Life	As supplied, 12 months from date of despatch
Acknowledgements	PrecisionAb™ is a trademark of Bio-Rad Laboratories
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: Antibody (10040): https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Rabbit IgG (H/L) (STAR208...) [HRP](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

'M295379:161101'

Printed on 30 Apr 2018