

Datasheet: VPA00269

Description:	RABBIT ANTI FGR
Specificity:	FGR
Format:	Purified
Product Type:	PrecisionAb™ Polyclonal
Isotype:	Polyclonal IgG
Quantity:	100 µl

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	■			1/1000

PrecisionAb antibodies have been extensively [validated for the western blot application](#). The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependant on sample type.

Target Species	Human
Product Form	Ig fraction - liquid
Preparation	Rabbit Ig fraction prepared by ammonium sulphate precipitation
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Immunogen	KLH-conjugated synthetic peptide corresponding to aa 3-33 of human FGR
External Database Links	UniProt: P09769 Related reagents Entrez Gene: 2268 FGR Related reagents
Synonyms	SRC2
Specificity	Rabbit anti Human FGR antibody recognizes FGR, also known as c-src-2 proto-oncogene, p55-c-fgr protein, proto-oncogene c-Fgr, tyrosine-protein kinase Fgr and v-fgr feline Gardner-Rasheed sarcoma viral oncogene homolog.

FGR is a member of the Src family of protein tyrosine kinases (PTKs). The FGR gene encoded protein contains N-terminal sites for myristylation and palmitoylation, a PTK domain, and SH2 and SH3 domains which are involved in mediating protein-protein interactions with phosphotyrosine-containing and proline-rich motifs, respectively. The protein localizes to plasma membrane ruffles, and functions as a negative regulator of cell migration and adhesion triggered by the beta-2 integrin signal transduction pathway. Infection with Epstein-Barr virus results in the overexpression of FGR. Multiple alternatively spliced variants, encoding the same protein, have been identified (provided by RefSeq, Jul 2008).

Rabbit anti Human FGR antibody detects a band of 56 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.

Western Blotting	Anti FGR detects a band of approximately 56 kDa in Ramos cell lysate
Instructions For Use	Please refer to the PrecisionAb western blotting protocol . For additional information on secondary antibody dilution and exposure time see product web page.
Storage	Store undiluted at -20°C, avoiding repeated freeze thaw cycles
Shelf Life	As supplied, 12 months from date of despatch
Acknowledgements	PrecisionAb™ is a trademark of Bio-Rad Laboratories
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: Antibody (10040): https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Rabbit IgG (H/L) (STAR208...) [HRP](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M286594:160718'

Printed on 30 Apr 2018