Datasheet: VPA00224

Description:	RABBIT ANTI TAK1
Specificity:	TAK1
Format:	Purified
Product Type:	PrecisionAb™ Polyclonal
Isotype:	Polyclonal IgG
Quantity:	100 µl

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	▪			1/1000

PrecisionAb antibodies have been extensively validated for the western blot application. The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependant on sample type.

Target Species

Human

Species Cross Reactivity

Reacts with: Mouse
N.B. Antibody reactivity and working conditions may vary between species.

Product Form

Purified IgG - liquid

Preparation

Rabbit polyclonal antibody purified by affinity chromatography on Protein G

Buffer Solution

Phosphate buffered saline

Preservative Stabilisers

0.09% Sodium Azide (NaN_3)

Immunogen

KLH conjugated synthetic peptide between 574-606 amino acids from the C-terminal region of human TAK1

External Database Links

UniProt:
O43318 Related reagents

Entrez Gene:
6885 MAP3K7 Related reagents

Synonyms

TAK1

Specificity	**Rabbit anti Human TAK1 antibody** recognizes TGF-beta activated kinase 1 (TAK1), also known as mitogen-activated protein kinase kinase kinase 7 and transforming growth factor-beta-activated kinase 1. Encoded by MAP3K7 gene, TAK1 is a member of the serine/threonine protein kinase family. This kinase mediates the signaling transduction induced by TGF beta and morphogenetic protein (BMP), and controls a variety of cell functions including transcription regulation and apoptosis. In response to IL-1, this protein forms a kinase complex including TRAF6, MAP3K7P1/TAB1 and MAP3K7P2/TAB2; this complex is required for the activation of nuclear factor kappa B. This kinase can also activate MAPK8/JNK, MAP2K4/MKK4, and thus plays a role in the cell response to environmental stresses. Four alternatively spliced transcript variants encoding distinct isoforms have been reported (provided by RefSeq, Jul 2008). Rabbit anti Human TAK1 antibody detects a band of 82 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.
Western Blotting	Anti TAK1 detects a band of approximately 82 kDa in HeLa cell lysates
Instructions For Use	Please refer to the PrecisionAb western blotting protocol. For additional information on secondary antibody dilution and exposure time see product web page.
Storage	Store undiluted at -20^{o}C, avoiding repeated freeze thaw cycles.
Shelf Life	As supplied, 12 months from date of despatch.
Acknowledgements	PrecisionAb™ is a trademark of Bio-Rad Laboratories.
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: Antibody (10040): https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Rabbit IgG (H/L) (STAR208...) HRP

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	**Worldwide**	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	**Europe**	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com

'M284930:160321'

Printed on 01 May 2018

© 2018 Bio-Rad Laboratories Inc | Legal | Imprint