

Datasheet: VPA00699

Description:	RABBIT ANTI XIAP
Specificity:	XIAP
Format:	Purified
Product Type:	PrecisionAb™ Polyclonal
Isotype:	Polyclonal IgG
Quantity:	100 µl

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	▪			1/1000

PrecisionAb antibodies have been extensively [validated for the western blot application](#). The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependant on sample type.

Target Species	Human
Species Cross Reactivity	Reacts with: Rat N.B. Antibody reactivity and working conditions may vary between species.
Product Form	Purified IgG - liquid
Preparation	Rabbit polyclonal antibody purified by affinity chromatography
Buffer Solution	TRIS buffered glycine
Preservative Stabilisers	0.01% Thiomersal <50% Glycerol
Immunogen	Recombinant protein encompassing part of the C-terminal region of human XIAP
External Database Links	<p>UniProt: P98170 Related reagents</p> <p>Entrez Gene: 331 XIAP Related reagents</p>
Synonyms	API3, BIRC4, IAP3

Specificity **Rabbit anti Human XIAP antibody** recognizes the E3 ubiquitin-protein ligase XIAP, also known as XIAP, XLP2, BIRC5 or MIHA.

The XIAP gene encodes a protein that belongs to a family of apoptotic suppressor proteins. Members of this family share a conserved motif termed, baculovirus IAP repeat, which is necessary for their anti-apoptotic function. This protein, BIRC5, functions through binding to tumor necrosis factor receptor-associated factors TRAF1 and TRAF2 and inhibits apoptosis induced by menadione, a potent inducer of free radicals, and interleukin 1-beta converting enzyme. MIHA also inhibits at least two members of the caspase family of cell-death proteases, caspase-3 and caspase-7. Mutations in XIAP are the cause of X-linked lymphoproliferative syndrome. Alternate splicing results in multiple transcript variants. Pseudogenes of XIAP are found on chromosomes 2 and 11 (provided by RefSeq, Feb 2011).

Rabbit anti Human XIAP antibody detects a band of 53 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.

Western Blotting Rabbit anti XIAP detects a band of approximately 53 kDa in MOLT-4 cell lysates

Instructions For Use Please refer to the [PrecisionAb western blotting protocol](#). For additional information on secondary antibody dilution and exposure time see product web page.

Storage Store undiluted at -20°C, avoiding repeated freeze thaw cycles.

Shelf Life As supplied, 12 months from date of despatch.

Acknowledgements PrecisionAb™ is a trademark of Bio-Rad Laboratories.

Health And Safety Information Material Safety Datasheet documentation #20357 available at: Antibody (20357): <https://www.bio-rad-antibodies.com/uploads/MSDS/20357.pdf>

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Rabbit IgG (H/L) (STAR208...) [HRP](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M302833:170201'

Printed on 02 May 2018